60 Rec'd PCT/PTO 04 DEC 2000

FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE ATTORNEY DOCKET NO. (REV 5-93) PATENT AND TRADEMARK OFFICE P108172-00022 DATE: December 4, 2000 TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) U.S. APPLAL NO.7 (IF KNOWN, SEE 37 C. CONCERNING A FILING UNDER 35 U.S.C. 371 Not yet assigned INTERNATIONAL APPLICATION NO. PRIORITY DATE CLAIMED INTERNATIONAL FILING DATE PCT/US99/12121 June 2, 1999 June 2, 1998 TITLE OF INVENTION: GENES OF CAROTENOID BIOSYNTHESIS AND METABOLISM AND METHODS OF USE THEREOF APPLICANT(S) FOR DO/EO/US: Francis CUNNINGHAM and Zairen SUN This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. (THE BASIC FILING FEE IS ATTACHED) This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. This express request to begin national examination procedures [35 U.S.C. 371(f)] at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). A proper demand for International Preliminary Amendment was made by the 19th month from the earliest claimed priority date. 4. A copy of the International Application as filed [35 U.S.C. 371(c)(2)] is transmitted herewith (required only if not transmitted by the International Bureau). а. 🔲 has been transmitted by the International Bureau. J is not required, as the application was filed in the United States Receiving Office (RO/US). 6. ☐ A translation of the International Application into English [35 U.S.C. 371(c)(2)]. Amendments to the claims of the International Application under PCT Article 19 [35 U.S.C. 371(c)(3)] 7. are transmitted herewith (required only if not transmitted by the International Bureau). ū have been transmitted by the International Bureau. In have not been made; however, the time limit for making such amendments has NOT expired. have not been made and will not be made. 33 8. A translation of the amendments to the claims under PCT Article 19 [35 U.S.C. 371(c)(3)]. 9. An oath or declaration of the inventor(s) [35 U.S.C. 371(c)(4)]. 10. A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 1 [35 U.S.C. 371(c)(5)]. Li Items 11 - 16 below concern other document(s) or information included: 11. An Information Disclosure Statement under 37 C.F.R. 1.97 and 1.98. 12.
An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. 3.28 and 3.31 is included. A FIRST preliminary amendment. ☐ A SECOND or SUBSEQUENT preliminary amendment. 14. A substitute specification. 15.
A change of power of attorney and/or address letter. 16. Other items or information: Copies of International Prel. Examination Reports (2); PCT/IB/308; PCT/IPEA/408; PCT/IB/332; Copy of Response to Written Opinion dated June 14, 2000; Statement; Computer readable form and paper copy of sequence listing; copy of published application (WO 99/63055) Drawings (45 sheets)

505 Rec'd PCT/PTO 04 DEC 2000

U.S. APPI N. NO. (IF KNOWN SEE 37 C.F.R. 1.50) Not yet assigned		INTERNATIONAL APPLICATION NO. PCT/US99/12121		ATTORNEY DOCKET NO. 108172-00022	
				DATE: December 4, 2000	
17. ☑ The following fees are submitted: Basic National Fee [37 C.F.R. 1.492(a)(1)-(5)]:				CALCULATIONS	PTO USE ONLY
Search Report has been prepared by the EPO or JPO\$860.00 International preliminary examination fee paid to USPTO (37 C.F.R. 1.482)\$690.00 No international preliminary examination fee paid to USPTO (37 C.F.R. 1.482) but international search fee paid to USPTO [37 C.F.R. 1.445(a)(2)]\$710.00 Neither international preliminary examination fee (37 C.F.R. 1.482) or international search fee [37 C.F.R. 1.445(a)(2)] paid to USPTO\$1,000.00 International preliminary examination fee paid to USPTO (37 C.F.R. 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)\$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 690.00	
Surcharge of \$130.00 for furnishing the path or declaration later				\$ 690.00	
than ☐ 20 ☐ 30 months from the earliest claimed priority date [37 C.F.R. 1.492(e)].					
Claims	Number Filed	Number Extra	Rate		
Total Claims	8 - 20 =	00	X \$ 18.00	\$	
Independent Claims	2 - 3 =	00	X \$ 80.00	\$	
Multiple dependent claim(s) (if applicable) + \$270.00				\$	
TOTAL OF ABOVE CALCULATIONS =				\$ 690.00	
Reduction by one-half for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 C.F.R. 1.9, 1.27, 1.28).				\$	
SUBTOTAL =				\$ 690.00	
Processing fee of \$130.00 for furnishing the English translation later the ☐ 20 ☐ 30 months from the earliest claimed priority date [37 C.F.R. 1.492(f)].				\$	
TOTAL NATIONAL FEE =				\$ 690.00	
Fee for recording the enclosed assignment [37 C.F.R. 1.21(h)]. The assignment must be accompanied by an appropriate cover sheet (37 C.F.R. 3.28, 3.31). \$40.00 per property +				\$	
TOTAL FEES ENCLOSED =				\$ 690.00	
				Amount to be refunded	\$
a. A check in the amount of \$ to cover the above fees is enclosed. b. Please charge my Deposit Account No. 01-2300 in the amount of \$690.00 to cover the above fee. A duplicate copy of this sheet is enclosed. c. A duplicate copy authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 01-2300. NOTE: Where an appropriate time limit under 37 C.F.R. 1.494 or 1.495 has not been met, a petition to revive [37 C.F.R. 1.137(a) or (b)] must be filed and granted to restore the application to pending status. SEND ALL CORRESPONDENCE TO: Arent Fox Kintner Plotkin & Kahn					
1050 Connecticut Avenue, N.W. Suite 600 Washington, D.C. 20036-5339 Tel: (202) 857-6000 Fax: (202) 638-4810 Richard J. Berman					

Rec'd PCT/PTO 18 JUN 2001 39/701395

PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:

CUNNINGHAM, Jr.

Appln. No.: 09/701,395

Filed: December 4, 2000 Attorney Dkt. No.: 108172-00022

For: GENES OF CAROTENOID BIOSYNTHESIS AND METABOLISM AND

METHODS OF USE THEREOF

PRELIMINARY AMENDMENT

Commissioner for Patents Washington, D.C. 20231

June 18, 2001

Sir:

Prior to [calculation of the filing fees and] initial examination of the application, please amend the above-identified application as follows:

IN THE SPECIFICATION:

Before Line 1, page 1 insert

-- CROSS-REFERENCE TO RELATED APPLICATION

This application is a National Stage entry of International Application No. PCT/US99/12121, filed June 2, 1999, the entire specification claims and drawings of which are incorporated herewith by reference. --

REMARKS

In the event that any fees are due with respect to the filing of this paper, please charge our Deposit Account No. 01-2300.

Respectfully submitted,

Richard J. Borman

Registration No. 39,107

ARENT FOX KINTNER PLOTKIN & KAHN, PLLC

Customer No. 004372

1050 Connecticut Avenue, N.W., Suite 600

Washington, D.C. 20036-5339

Tel: (202) 857-6000; Fax: (202) 638-4810

RJB/ccd

WO 99/63055

45/prds.

09/701395

PCT/US99/12121

525 Rec'd PCT/PTO 04 DEC 2000

GENES OF CAROTENOID BIOSYNTHESIS AND METABOLISM AND METHODS OF USE THEREOF

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention describes nucleic acid sequences for eukaryotic genes encoding ϵ lycopene ϵ -cyclase (also known as ϵ -cyclase and ϵ lycopene cyclase), isopentenyl pyrophosphate isomerase (IPP) and β -carotene hydroxylase as well as vectors containing the same and hosts transformed with said vectors. The present invention also provides methods for augmenting the accumulation of carotenoids, changing the composition of the carotenoids, and producing novel and rare carotenoids. The present invention provides methods for controlling the ratio or relative amounts of various carotenoids in a host. The invention also relates to modified lycopene ϵ -cyclase, IPP isomerase and β -carotene

hydroxylase. Additionally, the present invention provides a method for screening for genes and cDNAs encoding enzymes of carotenoid biosynthesis and metabolism.

Background of the Invention

Carotenoid pigments with cyclic endgroups are essential components of the photosynthetic apparatus in oxygenic photosynthetic organisms (e.g., cyanobacteria, algae and plants; Goodwin, 1980). The symmetrical bicyclic yellow carotenoid pigment βcarotene (or, in rare cases, the asymmetrical bicyclic α -carotene) is intimately associated with the photosynthetic reaction centers and plays a vital role in protecting against potentially lethal photooxidative damage (Koyama, 1991). β-carotene and other carotenoids derived from it or from α-carotene also serve as light-harvesting pigments (Siefermann-Harms, 1987), are involved in the thermal dissipation of excess light energy captured by the lightharvesting antenna (Demmig-Adams & Adams, 1992), provide substrate for the biosynthesis of the plant growth regulator abscisic acid (Rock & Zeevaart, 1991; Parry & Horgan, 1991), and are precursors of vitamin A in human and animal diets (Krinsky, 1987). Plants also exploit carotenoids as coloring agents in flowers and fruits to attract pollinators and agents of seed dispersal (Goodwin, 1980). The color provided by carotenoids is also of agronomic value in a number of important crops. Carotenoids are currently harvested from a variety of organisms, including plants, algae, yeasts, cyanobacteria and bacteria, for use as pigments in food and feed.

5

10

25

30

30

5

10

The probable pathway for formation of cyclic carotenoids in plants, algae and cyanobacteria is illustrated in Figure 1. Two types of cyclic endgroups or rings are commonly found in higher plant carotenoids, these are referred to as the β (beta) and ϵ (epsilon) rings (Fig. 3). The precursor acyclic endgroup (no ring structure) is referred to as the Ψ (psi) endgroup. The β and ϵ endgroups differ only in the position of the double bond in the ring. Carotenoids with two β rings are ubiquitous, and those with one β and one ϵ ring are common, but carotenoids with two ϵ rings are uncommon. β -carotene (Fig. 1) has two β -endgroups and is a symmetrical compound that is the precursor of a number of other important plant carotenoids such as zeaxanthin and violaxanthin (Fig. 2).

Genes encoding enzymes of carotenoid biosynthesis have previously been isolated from a variety of sources including bacteria (Armstrong et al., 1989, Mol. Gen. Genet. 216, 254-268; Misawa et al., 1990, J. Bacteriol., 172, 6704-12), fungi (Schmidhauser et al., 1990, Mol. Cell. Biol. 10, 5064-70), cyanobacteria (Chamovitz et al., 1990, Z. Naturforsch, 45c, 482-86; Cunningham et al., 1994) and higher plants (Bartley et al., Proc. Natl. Acad. Sci USA 88, 6532-36; Martinez-Ferez & Vioque, 1992, Plant Mol. Biol. 18, 981-83). Many of the isolated enzymes show a great diversity in structure, function and inhibitory properties between sources. For example, phytoene desaturases from the cyanobacterium Synechococcus and from higher plants and green algae carry out a two-step desaturation to yield ζ-carotene as a reaction product. In plants and cyanobacteria a second enzyme (ζcarotene desaturase), similar in amino acid sequence to the phytoene desaturase, catalyzes two additional desaturations to yield lycopene. In contrast, a single desaturase enzyme from Erwinia herbicola and from other bacteria introduces all four double bonds required to form lycopene. The Erwinia and other bacterial desaturases bear little amino acid sequence similarity to the plant and cyanobacterial desaturase enzymes, and are thought to be of unrelated ancestry. Therefore, even with a gene in hand from one source, it may be difficult to identify a gene encoding an enzyme of similar function in another organism. In particular, the sequence similarity between certain of the prokaryotic and eukaryotic genes encoding enzymes of carotenoid biosynthesis is quite low.

Further, the mechanism of gene expression in prokaryotes and eukaryotes appears to differ sufficiently such that one cannot expect that an isolated eukaryotic gene will be properly expressed in a prokaryotic host.

30

5

10

The difficulties in isolating genes encoding enzymes with similar functions is exemplified by recent efforts to isolate the gene encoding the enzyme that catalyzes the formation of β -carotene from the acyclic precursor lycopene. Although a gene encoding an enzyme with this function had been isolated from a bacterium, it had not been isolated from any photosynthetic procaryote or from any eukaryotic organism. The isolation and characterization of the enzyme catalyzing formation of β -carotene in the cyanobacterium Synechococcus PCC7942 was described by the present inventors and others (Cunningham et al., 1993 and 1994). The amino acid sequence similarity of the cyanobacterial enzyme to the various bacterial lycopene β -cyclases is so low (ca. 18-25% overall; Cunningham et al., 1994) that there is much uncertainty as to whether they share a common ancestry or, instead, represent an example of convergent evolution.

The need remains for the isolation of eukaryotic and prokaryotic genes and cDNAs encoding polypeptides involved in the carotenoid biosynthetic pathway, including those encoding a lycopene ϵ -cyclase, IPP isomerase and β -carotene hydroxylase. There remains a need for methods to enhance the production of carotenoids, to alter the composition of carotenoids, and to reduce or eliminate carotenoid production. There also remains a need in the art for methods for screening for genes and cDNAs encoding enzymes of carotenoid biosynthesis and metabolism.

SUMMARY OF THE INVENTION

Accordingly, a first object of this invention is to provide purified and/or isolated nucleic acids which encode enzymes involved in carotenoid biosynthesis; in particular, lycopene ϵ -cyclase, IPP isomerase and β -carotene hydroxylase.

A second object of this invention is to provide purified and/or isolated nucleic acids which encode enzymes which produce novel or uncommon carotenoids.

A third object of the present invention is to provide vectors containing said genes.

A fourth object of the present invention is to provide hosts transformed with said vectors.

Another object of the present invention is to provide hosts which accumulate novel or uncommon carotenoids or which accumulate greater amounts of specific or total carotenoids.

Another object of the present invention is to provide hosts with inhibited and/or altered carotenoid production.

10

Another object of this invention is to secure the expression of eukaryotic carotenoidrelated genes in a recombinant prokaryotic host.

Yet another object of the present invention is to provide a method for screening for eukaryotic and prokaryotic genes and cDNAs which encode enzymes involved in carotenoid biosynthesis and metabolism.

An additional object of the invention is to provide a method for manipulating carotenoid biosynthesis in photosynthetic organisms by inhibiting the synthesis of certain enzymatic products to cause accumulation of precursor compounds.

Another object of the invention is to provide modified lycopene ϵ -cyclase, IPP isomerase and β -carotene hydroxylase.

These and other objects of the present invention have been realized by the present inventors as described below.

A subject of the present invention is an isolated and/or purified nucleic acid sequence which encodes for a protein having lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase enzyme activity and having the amino acid sequence of SEQ ID NOS: 2, 4, 14-21 or 23-27.

The invention also includes vectors which comprise any of the nucleic acid sequences listed above, and host cells transformed with such vectors.

Another subject of the present invention is a method of producing or enhancing the production of a carotenoid in a host cell, comprising inserting into the host cell a vector comprising a heterologous nucleic acid sequence which encodes for a protein having lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase enzyme activity, wherein the heterologous nucleic acid sequence is operably linked to a promoter; and expressing the heterologous nucleic acid sequence to produce the protein.

Yet another subject of the present invention is a method of modifying the production of carotenoids in a host cell, the method comprising inserting into the host cell a vector comprising a heterologous nucleic acid sequence which produces an RNA and/or encodes for a protein which modifies lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase enzyme activity, relative to an untransformed host cell, wherein the heterologous nucleic acid sequence is operably linked to a promoter; and expressing the heterologous nucleic acid sequence in the host cell to modify the production of the carotenoids in the host cell, relative to the untransformed host cell.

30

25

WO 99/63055

5

10

The present invention also includes a method of expressing, in a host cell, a heterologous nucleic acid sequence which encodes for a protein having lycopene ε-cyclase, IPP isomerase or β-carotene hydroxylase enzyme activity, the method comprising inserting into the host cell a vector comprising the heterologous nucleic acid sequence, wherein the heterologous nucleic acid sequence is operably linked to a promoter; and expressing the heterologous nucleic acid sequence.

Also included is a method of expressing, in a host cell, a heterologous nucleic acid sequence which encodes for a protein which modifies lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase enzyme activity in the host cell, relative to an untransformed host cell, the method comprising inserting into the host cell a vector comprising the heterologous nucleic acid sequence, wherein the heterologous nucleic acid sequence is operably linked to a promoter; and expressing the heterologous nucleic acid sequence.

Another subject of the present invention is a method for screening for genes and cDNAs which encode enzymes involved in carotenoid biosynthesis and metabolism.

BRIEF DESCRIPTION OF THE DRAWINGS

A more complete appreciation of the invention and many of the attendant advantages thereof will be readily obtained as the same becomes better understood by reference to the following detailed description when considered in connection with the accompanying drawings, wherein:

Figure 1 is a schematic representation of the putative pathway of β -carotene biosynthesis in cyanobacteria, algae and plants. The enzymes catalyzing various steps are indicated at the left. Target sites of the bleaching herbicides NFZ and MPTA are also indicated at the left. Abbreviations: DMAPP, dimethylallyl pyrophosphate; FPP, farnesyl pyrophosphate; GGPP, geranylgeranyl pyrophosphate; GPP, geranyl pyrophosphate; IPP, isopentenyl pyrophosphate; LCY, lycopene cyclase; MVA, mevalonic acid; MPTA, 2-(4-methylphenoxy)triethylamine hydrochloride; NFZ, norflurazon; PDS, phytoene desaturase; PSY, phytoene synthase; ZDS, ζ -carotene desaturase; PPPP, prephytoene pyrophosphate.

Figure 2 depicts possible routes of synthesis of cyclic carotenoids and common plant and algal xanthophylls (oxycarotenoids) from neurosporene. Demonstrated activities of the β - and ϵ -cyclase enzymes of A. thaliana are indicated by bold arrows labelled with β or ϵ respectively. A bar below the arrow leading to ϵ -carotene indicates that the enzymatic

25

30

5

10

activity was examined but no product was detected. The steps marked by an arrow with a dotted line have not been specifically examined. Conventional numbering of the carbon atoms is given for neurosporene and α -carotene. Inverted triangles (∇) mark positions of the double bonds introduced as a consequence of the desaturation reactions.

Figure 3 depicts the carotene endgroups which are found in plants.

Figure 4 is a DNA sequence and the predicted amino acid sequence of a lycopene ϵ -cyclase cDNA isolated from A. thaliana (SEQ ID NOS: 1 and 2). These sequences were deposited under Genbank accession number U50738. This cDNA is incorporated into the plasmid pATeps.

Figure 5 is a DNA sequence encoding the β -carotene hydroxylase isolated from A. thaliana (SEQ ID NO: 3). This cDNA is incorporated into the plasmid pATOHB.

Figure 6 is an alignment of the predicted amino acid sequences of *A. thaliana* β-carotene hydroxylase (SEQ ID NO: 4) with those of the bacterial β-carotene hydroxylase enzymes from *Alicalgenes sp.* (SEQ ID NO: 5) (Genbank D58422), *Erwinia herbicola* Eho10 (SEQ ID NO.: 6) (GenBank M872280), *Erwinia uredovora* (SEQ ID NO.: 7) (GenBank D90087) and *Agrobacterium aurianticum* (SEQ ID NO.: 8) (GenBank D58420). A consensus sequence is also shown. All five genes are identical where a capital letter appears in the consensus. A lowercase letter indicates that three of five, including *A. thaliana*, have the identical residue. TM; transmembrane.

Figure 7 is a DNA sequence of a cDNA encoding an IPP isomerase isolated from A. thaliana (SEQ ID NO: 9). This cDNA is incorporated into the plasmid pATDP5.

Figure 8 is a DNA sequence of a second cDNA encoding another IPP isomerase isolated from A. thaliana (SEQ ID NO: 10). This cDNA is incorporated into the plasmid pATDP7.

Figure 9 is a DNA sequence of a cDNA encoding an IPP isomerase isolated from *Haematococcus pluvialis* (SEQ ID NO: 11). This cDNA is incorporated into the plasmid pHP04.

Figure 10 is a DNA sequence of a second cDNA encoding another IPP isomerase isolated from *Haematococcus pluvialis* (SEQ ID NO: 12). This cDNA is incorporated into the plasmid pHP05.

Figure 11 is an alignment of the amino acid sequences predicted by IPP isomerase cDNAs isolated from A. thaliana (SEQ ID NO.: 16 and 18), H. pluvialis (SEQ ID NOS.: 14

30

5

10

and 15), Clarkia breweri (SEQ ID NO.: 17) (See, Blanc & Pichersky, Plant Physiol. (1995) 108:855; Genbank accession no. X82627) and Saccharomyces cerevisiae (SEQ ID NO.: 19) (Genbank accession no. J05090).

Figure 12 is a DNA sequence of the cDNA encoding an IPP isomerase isolated from *Tagetes erecta* (marigold; SEQ ID NO: 13). This cDNA is incorporated into the plasmid pPMDP1. xxx's denote a region not originally sequenced. Figure 21A shows the complete marigold sequence.

Figure 13 is an alignment of the consensus sequence of four plant β -cyclases (SEQ ID NO.: 20) with the A. thaliana lycopene ϵ -cyclase (SEQ ID NO.: 21). A capital letter in the plant β consensus is used where all four β -cyclase genes predict the same amino acid residue in this position. A small letter indicates that an identical residue was found in three of the four. Dashes indicate that the amino acid residue was not conserved and dots in the sequence denote a gap. A consensus for the aligned sequences is given, in capital letters below the alignment, where the β - and ϵ -cyclases have the same amino acid residue. Arrows indicate some of the conserved amino acids that will be used as junction sites for construction of chimeric cyclases with novel enzymatic activities. Several regions of interest including a sequence signature indicative of a dinucleotide-binding motif and two predicted transmembrane (TM) helical regions are indicated below the alignment and are underlined.

Figure 14 shows the nucleotide (SEQ ID NO:22) and amino acid sequences (SEQ ID NO:23) of the *Adonis palaestina* (pheasant's eye) ∈-cyclase cDNA #5.

Figure 15A shows the nucleotide (SEQ ID NO:24) and amino acid sequences (SEQ ID NO:25) of a potato ε-cyclase cDNA. Figure 15B shows the amino acid sequence (SEQ ID NO:26) of a chimeric lettuce/potato lycopene ε-cyclase. Amino acids in lower case are from the lettuce cDNA and those in upper case are from the potato cDNA. The product of this chimeric cDNA has e-cyclase activity and converts lycopene to the monocyclic δ-carotene.

Figure 16 shows a comparison between the amino acid sequences of the *Arabidopsis* ϵ -cyclase (SEQ ID NO:27) and the potato ϵ -cyclase (SEQ ID NO:25).

Figure 17A shows the nucleotide sequence of the *Adonis palaestina* Ipi1 (SEQ ID NO:28) and Figure 17B shows the nucleotide sequence of the *Adonis palaestina* Ipi2 (SEQ ID NO: 29).

30

5

10

Figure 18A shows the nucleotide sequence of the *Haematoccus pluvialis* Ipi1 (SEQ ID NO:11) and Figure 18B shows the nucleotide sequence of the *Haematoccus pluvialis* Ipi2 (SEQ ID NO:30).

Figure 19A shows the nucleotide sequence of the *Lactuca sativa (romaine lettuce)*Ipi1 (SEQ ID NO:31) and Figure 19B shows the nucleotide sequence of the *Lactuca sativa*Ipi2 (SEQ ID NO: 32).

Figure 20 shows the nucleotide sequence of the *Chlamydomonas reinhardtii* Ipi1 (SEO ID NO:33).

Figure 21A shows the nucleotide sequence of the *Tagetes erecta* (marigold) Ipi1 (SEQ ID NO:34) and Figure 21B shows the nucleotide sequence of the *Oryza sativa* (rice) Ipi1 (SEQ ID NO:35).

Figure 22 shows a amino acid sequence alignment of various plant and green algal isopentenyl isomerases (IPI) (SEQ ID NOS:16, 36-45).

Figure 23 shows a comparison between *Adonis palaestina* ϵ -cyclase cDNA #3 and cDNA #5 nucleotide sequences.

Figure 24 shows a comparison between *Adonis palaestina* ϵ -cyclase cDNA #3 and cDNA #5 predicted amino acid sequences.

Figure 25 shows a sequence alignment of various plant β - and ϵ -cyclases. Those sequences outlined in grey denote identical sequences among the ϵ -cyclases. Those sequences outlined in black denote identical sequences among both the β - and ϵ -cyclases.

Figure 26 shows a sequence alignment of the plant ϵ -cyclases from Figure 25. Those sequences outlined in black denote identical sequences among the ϵ -cyclases.

Figure 27 is a dendrogram or "tree" illustrating the degree of amino acid sequence similarity for various lycopene β - and ϵ -cyclases.

Figure 28 shows a comparison between Arabidopsis ϵ -cyclase and lettuce ϵ -cyclase predicted amino acid sequences.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention includes an isolated and/or purified nucleic acid sequence which encodes for a protein having lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase enzyme activity and having the amino acid sequence of SEQ ID NOS: 2, 4, 14-21, 23 or 25-27. Nucleic acids encoding lycopene ϵ -cyclase, β -carotene hydroxylase and IPP

30

5

10

isomerases have been isolated from several genetically distant sources.

The present inventors have isolated nucleic acids encoding the enzyme IPP isomerase, which catalyzes the reversible conversion of isopentenyl pyrophosphate (IPP) to dimethylallyl pyrophosphate (DMAPP). IPP isomerase cDNAs were isolated from the plants A. thaliana, Tagetes erecta (marigold), Adonis palaestina (pheasant's eye), Lactuca sativa (romaine lettuce) and from the green algae H. pluvialis and Chlamydomonas reinhardtii. Alignments of the amino acid sequences predicted by some of these cDNAs are shown in Figures 12 and 22. Plasmids containing some of these cDNAs were deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville MD 20852 on March 4, 1996 under ATCC accession numbers 98000 (pHP05 - H. pluvialis); 98001 (pMDP1 - marigold); 98002 (pATDP7 - A. thaliana) and 98004 (pHP04 - H. pluvialis).

The present inventors have also isolated nucleic acids encoding the enzyme β -carotene hydroxylase, which is responsible for hydroxylating the β -endgroup in carotenoids. The nucleic acid of the present invention is shown in SEQ ID NO: 3 and Figure 5. The full length cDNA product hydroxylates both end groups of β -carotene as do products of cDNAs which encode proteins truncated by up to 50 amino acids from the N-terminus. Products of genes which encode proteins truncated between about 60-110 amino acids from the N-terminus preferentially hydroxylate only one ring. A plasmid containing this gene was deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville MD 20852 on March 4, 1996 under ATCC accession number 98003 (pATOHB - *A. thaliana*).

The present inventors have also isolated nucleic acids encoding the enzyme lycopene ϵ -cyclase, which is responsible for the formation of ϵ -endgroups in carotenoids. The A. thaliane ϵ -cyclase adds an ϵ ring to only one end of the symmetrical lycopene while the related β -cyclase adds a ring at both ends. The A. thaliana cDNA of the present invention is shown in Figure 4 and SEQ ID NO: 1. A plasmid containing this gene was deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville MD 20852 on March 4, 1996 under ATCC accession number 98005 (pATeps - A. thaliana).

In addition, lycopene ϵ -cyclases have been identified in lettuce and in *Adonis* palaestina (cDNA #5) which encode enzymes that convert lycopene to the bicyclic ϵ -carotene (ϵ , ϵ -carotene). An additional cDNA from *Adonis palaestina* (cDNA #3) encodes a lycopene ϵ -cyclase which converts lycopene into δ -carotene (ϵ , ψ -carotene) and differs from the lycopene ϵ -cyclase which forms bicyclic ϵ -carotene (ϵ , ϵ -carotene) by only 5 amino acids.

30

5

10

One or more of these amino acids may be modified by alteration of the nucleotide sequence in the #5 cDNA to obtain an enzyme which forms the bicyclic ϵ, ϵ -carotene. The sequences of the *Adonis palaestina* and *Arabidopsis thaliana* ϵ -cyclases have about 70% nucleotide identity and about 72% amino acid identity.

identity and about 72% amino acid identity.

Initial experiments by the inventors with chimeric genes indicated that the part of the

∈-cyclase which is responsible for adding 2 ∈ rings to form ∈,∈-carotene is the carboxy
terminal portion of the gene. The lettuce ∈-cyclase adds two ∈ rings to form ∈,∈-carotene. A

DNA encoding a partial potato ϵ -cyclase (missing its amino terminal portion), when combined with an amino terminal region from the lettuce ϵ -cyclase gene, produces a

monocyclic δ -carotene (ϵ , ψ -carotene). With the discovery of the differences between the

Adonis palaestina clone #3 and clone #5, the specific amino acids responsible for the addition

of an extra ϵ ring have been identified (Figure 24). Specifically, amino acid 55 is Thr in

clone #3 and Ser in clone #5, amino acid 210 is Asn in clone #3 and Asp in clone #5, amino

acid 231 is Asp in clone #3 and Glu in clone #5, amino acid 352 is Ile in clone #3 and Val in

clone #5, and amino acid 524 is Lys in clone #3 and Arg in clone #5. It can be appreciated

that these changes are quite conservative, as only one change, at amino acid 210, changes the

charge of the protein.

Thus, it is clear that the nucleic acids of the invention encoding the enzymes as presently disclosed may be altered to increase a particularly desirable property of the enzyme, to change a property of the enzyme, or to diminish an undesirable property of the enzyme. Such modifications can be by deletion, substitution, or insertion of one or more amino acids, and can be performed by routine enzymatic manipulation of the nucleic acid encoding the enzyme (such as by restriction enzyme digestion, removal of nucleotides by mung bean nuclease or *Bal31*, insertion of nucleotides by Klenow fragment, and by religation of the ends), by site-directed mutagenesis, or may be accidental, such as by low fidelity PCR or those obtained through mutations in hosts that are producers of the enzymes. These techniques as well as other suitable techniques are well known in the art.

Mutations can be made in the nucleic acids of the invention such that a particular codon is changed to a codon which codes for a different amino acid. Such a mutation is generally made by making the fewest nucleotide changes possible. A substitution mutation of this sort can be made to change an amino acid in the resulting protein in a non-conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping

30

5

of amino acids having a particular size or characteristic to an amino acid belonging to another grouping) or in a conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping of amino acids having a particular size or characteristic to an amino acid belonging to the same grouping). Such a conservative change generally leads to less change in the structure and function of the resulting protein. A non-conservative change is more likely to alter the structure, activity or function of the resulting protein. The present invention should be considered to include sequences containing conservative changes which do not significantly alter the activity or binding characteristics of the resulting protein.

The following is one example of various groupings of amino acids:

Amino acids with nonpolar R groups: Alanine, Valine, Leucine, Isoleucine, Proline, Phenylalanine, Tryptophan and Methionine.

Amino acids with uncharged polar R groups: Glycine, Serine, Threonine, Cysteine, Tyrosine, Asparagine and Glutamine.

Amino acids with charged polar R groups (negatively charged at Ph 6.0): Aspartic acid and Glutamic acid.

Basic amino acids (positively charged at pH 6.0): Lysine, Arginine and Histidine.

Another grouping may be those amino acids with phenyl groups: Phenylalanine, Tryptophan and Tyrosine.

Another grouping may be according to molecular weight (i.e., size of R groups). Particularly preferred substitutions are:

- Lys for Arg and vice versa such that a positive charge may be maintained;
- Glu for Asp and vice versa such that a negative charge may be maintained;
- Ser for Thr such that a free -OH can be maintained; and
- Gln for Asn such that a free NH₂ can be maintained.

Amino acid substitutions may also be introduced to substitute an amino acid with a particularly preferable property. For example, a Cys may be introduced to provide a potential site for disulfide bridges with another Cys. A His may be introduced as a particularly "catalytic" site (i.e., His can act as an acid or base and is the most common amino acid in biochemical catalysis). Pro may be introduced because of its particularly planar structure, which induces β-turns in the protein's structure.

It is clear that certain modifications of SEQ ID NOS: 2, 4, 14-21, 23 or 25-27 can take place without destroying the activity of the enzyme. It is noted especially that truncated

30

5

10

versions of the nucleic acids of the invention are functional. For example, several amino acids (from 1 to about 120) can be deleted from the N-terminus of the lycopene ϵ -cyclases of the invention, and a functional protein can still be produced. This fact is made especially clear from Figure 25, which shows a sequence alignment of several plant ϵ -cyclases. As can be seen from Figure 25, there is an enormous amount of sequence disparity between amino acid sequences 2 to about 50-70 (depending on the particular sequence, since gaps are present). There is less, but also a substantial amount of, sequence dissimilarity between about 50-70 to about 90-120 (depending on the particular sequence). Thereafter, the sequences are fairly conserved, except for small pockets of dissimilarity between about 275-295 to about 285-305 (depending on the particular sequence), and between about 395-415 to about 410-430 (depending on the particular sequence).

The present inventors have found that the amount of the 5' region present in the nucleic acids of the invention can alter the activity of the enzyme. Instead of diminishing activity, truncating the 5' region of the nucleic acids of the invention may result in an enzyme with a different specificity. Thus, the present invention relates to nucleic acids and enzymes encoded thereby which are truncated to within 0-50, preferably 0-25, codons of the 5' initiation codon of their prokaryotic counterparts as determined by alignment maps as discussed below.

For example, when the cDNA encoding A. thaliana β -carotene hydroxylase was truncated, the resulting enzyme catalyzed the formation of β -cryptoxanthin as the major product and zeaxanthin as minor product; in contrast to its normal production of zeaxanthin.

The present invention is intended to include those nucleic acid and amino acid sequences in which substitutions, deletions, additions or other modifications have taken place, as compared to SEQ ID NOS: 2, 4, 14-21, 23 or 25-27, without destroying the activity of the enzyme. Preferably, the substitutions, deletions, additions or other modifications take place at the 5' end, or any other of those positions which already show dissimilarity between any of the presently disclosed amino acid sequences (see also Figure 25) or other amino acid sequences which are known in the art and which encode the same enzyme (i.e., lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase).

In each case, nucleic acid and amino acid sequence similarity and identity is measured using sequence analysis software, for example, the Sequence Analysis, Gap, or BestFit software packages of the Genetics Computer Group (University of Wisconsin Biotechnology

30

5

10

Center, 1710 University Avenue, Madison, Wisconsin 53705), MEGAlign (DNAStar, Inc., 1228 S. Park St., Madison, Wisconsin 53715), or MacVector (Oxford Molecular Group, 2105 S. Bascom Avenue, Suite 200, Campbell, California 95008). Such software uses algorithms to match similar sequences by assigning degrees of identity to various substitutions, deletions, and other modifications, and includes detailed instructions as to useful parameters, etc., such that those of routine skill in the art can easily compare sequence similarities and identities. An example of a useful algorithm in this regard is the algorithm of Needleman and Wunsch, which is used in the Gap program discussed above. This program finds the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. Another useful algorithm is the algorithm of Smith and Waterman, which is used in the BestFit program discussed above. This program creates an optimal alignment of the best segment of similarity between two sequences. Optimal alignments are found by inserting gaps to maximize the number of matches using the local homology algorithm of Smith and Waterman.

Conservative (i.e. similar) substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine and leucine; aspartic acid, glutamic acid, asparagine and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine. Substitutions may also be made on the basis of conserved hydrophobicity or hydrophilicity (see Kyte and Doolittle, *J. Mol. Biol.* 157: 105-132 (1982)), or on the basis of the ability to assume similar polypeptide secondary structure (see Chou and Fasman, *Adv. Enzymol.* 47: 45-148 (1978)).

If comparison is made between nucleotide sequences, preferably the length of comparison sequences is at least 50 nucleotides, more preferably at least 60 nucleotides, at least 75 nucleotides or at least 100 nucleotides. It is most preferred if comparison is made between the nucleic acid sequences encoding the enzyme coding regions necessary for enzyme activity. If comparison is made between amino acid sequences, preferably the length of comparison is at least 20 amino acids, more preferably at least 30 amino acids, at least 40 amino acids or at least 50 amino acids. It is most preferred if comparison is made between the amino acid sequences in the enzyme coding regions necessary for enzyme activity.

It should be appreciated that also within the scope of the present invention are nucleic acid sequences encoding lycopene ϵ -cyclases, IPP isomerases and β -carotene hydroxylases

30

5

10

which code for enzymes having the same amino acid sequence as SEQ ID NOS: 2, 4, 14-21, 23 or 25-27, but which are degenerate to the nucleic acids specifically disclosed herein.

The amino acid residues described herein are preferred to be in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property of immunoglobulin-binding is retained by the polypeptide.

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook et al, "Molecular Cloning: A Laboratory Manual" (1989); "Current Protocols in Molecular Biology" Volumes I-III [Ausubel, R. M., ed. (1994)]; "Cell Biology: A Laboratory Handbook" Volumes I-III [J. E. Celis, ed. (1994))]; "Current Protocols in Immunology" Volumes I-III [Coligan, J. E., ed. (1994)]; "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" [B.D. Hames & S.J. Higgins eds. (1985)]; "Transcription And Translation" [B.D. Hames & S.J. Higgins, eds. (1984)]; "Animal Cell Culture" [R.I. Freshney, ed. (1986)]; "Immobilized Cells And Enzymes" [IRL Press, (1986)]; B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

The present invention also includes vectors. Suitable vectors according to the present invention comprise a nucleic acid of the invention encoding an enzyme involved in carotenoid biosynthesis or metabolism and a suitable promoter for the host, and can be constructed using techniques well known in the art (for example Sambrook et al., Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989; Ausubel et al., Current Protocols in Molecular Biology, Greene Publishing and Wiley Interscience, New York, 1991). Suitable vectors for eukaryotic expression in plants are described in Frey et al., Plant J. (1995) 8(5):693 and Misawa et al, 1994a; incorporated herein by reference. Suitable vectors for prokaryotic expression include pACYC184, pUC119, and pBR322 (available from New England BioLabs, Bevery, MA) and pTrcHis (Invitrogen) and pET28 (Novagen) and derivatives thereof. The vectors of the present invention can additionally contain regulatory elements such as promoters, repressors, selectable markers such as antibiotic resistance genes, etc.

The nucleic acids encoding the carotenoid enzymes as described above, when cloned into a suitable expression vector, can be used to overexpress these enzymes in a plant

10

30

25

expression system or to inhibit the expression of these enzymes. For example, a vector containing the gene encoding lycopene ϵ -cyclase can be used to increase the amount of α -carotene and carotenoids derived from α -carotene (such as lutein and α -cryptoxanthin) in an organism and thereby alter the nutritional value, pharmacology and visual appearance value of the organism.

Therefore, the present invention includes a method of producing or enhancing the production of a carotenoid in a host cell, relative to an untransformed host cell, the method comprising inserting into the host cell a vector comprising a heterologous nucleic acid sequence which encodes for a protein having lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase enzyme activity, wherein the heterologous nucleic acid sequence is operably linked to a promoter; and expressing the heterologous nucleic acid sequence to produce the protein.

The present invention also includes a method of modifying the production of carotenoids in a host cell, the method comprising inserting into the host cell a vector comprising a heterologous nucleic acid sequence which produces an RNA and/or encodes for a protein which modifies lycopene ϵ -cyclase, IPP isomerase or β -carotene hydroxylase enzyme activity, relative to an untransformed host cell, wherein the heterologous nucleic acid sequence is operably linked to a promoter; and expressing the heterologous nucleic acid sequence in the host cell to modify the production of the carotenoids in the host cell, relative to the untransformed host cell.

The term "modifying the production" means that the amount of carotenoids produced in the host cell can be enhanced, reduced, or left the same, as compared to the untransformed host cell. In accordance with one embodiment of the present invention, the make-up of the carotenoids (i.e., the specific carotenoids produced) is changed vis a vis each other, and this change in make-up may result in either a net gain, net loss, or no net change in the total amount of carotenoids produced in the cell. In accordance with another embodiment of the present invention, the production or the biochemical activity of the carotenoids (or the enzymes which catalyze their formation) is enhanced by the insertion of an enzyme-encoding nucleic acid of the invention. In yet another embodiment of the invention, the production or the biochemical activity of the carotenoids (or the enzymes which catalyze their formation) may be reduced or inhibited by a number of different approaches available to those skilled in the art, including but not limited to such methodologies or approaches as anti-sense (e.g.,

30

5

10

Gray et al (1992) Plant Mol. Biol. 19:69-87), ribozymes (e.g., Wegener et al (1994) Mol. Gen. Genet. 245:465-470), co-suppression (e.g., Fray and Grierson (1993) Plant Mol. Biol. 22:589-602), targeted disruption of the gene (e.g., Schaefer et al. (1997) Plant J. 11:1195-1206), intracellular antibodies (e.g., Rondon and Marasco (1997) Ann. Rev. Microbiol. 51:257-283) or whatever other approaches rely on the knowledge or availability of the nucleic acid or amino acid sequences of the invention and/or portions thereof, to thereby reduce accumulation of carotenoids with ϵ rings and compounds derived from them (for ϵ -cyclase inhibition), or carotenoids with hydroxylated β rings and compounds derived from them (for β-hydroxylase inhibition), or, in the case if IPP isomerase, accumulation of any isoprenoid compound.

Preferably, at least a portion of the nucleic acid sequences used in the methods, vectors and host cells of the invention codes for an enzyme having an amino acid sequence which is at least 85% identical, preferably at least 90%, at least 95% or completely identical to SEQ ID NOS: 2, 4, 14-21, 23 or 25-27. Sequence identity is determined as noted above. Preferably, sequence additions, deletions or other modifications are made as indicated above. so as to not affect the function of the particular enzyme.

In a preferred embodiment, vectors are manufactured which contain a DNA encoding a eukaryotic IPP isomerase upstream of a DNA encoding a second eukaryotic carotenoid enzyme. The inventors have discovered that inclusion of an IPP isomerase gene increases the supply of substrate for the carotenoid pathway; thereby enhancing the production of carotenoid endproducts, as compared to a host cell which is not transformed with such a vector. This is apparent from the much deeper pigmentation in carotenoid-accumulating colonies of E. coli which also contain one of the aforementioned IPP isomerase genes when compared to colonies that lack this additional IPP isomerase gene. Similarly, a vector comprising an IPP isomerase gene can be used to enhance production of any secondary metabolite of dimethylallyl pyrophosphate and/or isopentenyl pyrophosphate (such as isoprenoids, steroids, carotenoids, etc.). The term "isoprenoid" is intended to mean any member of the class of naturally occurring compounds whose carbon skeletons are composed. in part or entirely, of isopentyl C, units. Preferably, the carbon skeleton is of an essential oil, a fragrance, a rubber, a carotenoid, or a therapeutic compound, such as paclitaxel.

A vertor containing the cDNA encoding a lycopene ϵ -cyclase of the invention. preferably the lettuce lycopene ϵ -cyclase or Adonis ϵ -cyclase #5, can be used to increase the

10

25

30

amount of bicyclic €-carotene in an organism and thereby alter the nutritional value, pharmacology and visual appearance value of the organism. In addition, the transformed organism can be used in the formulation of therapeutic agents, for example in the treatment of cancer (see Mayne et al (1996) FASEB J. 10:690-701; Tsushima et al (1995) Biol. Pharm. Bull. 18:227-233).

An antisense strand of a nucleic acid of the invention can be inserted into a vector. For example, the lycopene ϵ -cyclase gene can be inserted into a vector and incorporated into the genomic DNA of a host, thereby inhibiting the synthesis of ϵ , β -carotenoids (lutein and α -carotene) and enhancing the synthesis of β , β -carotenoids (zeaxanthin and β -carotene).

The present invention also relates to novel enzymes which are encoded by the amino acid sequences of the invention, or portions thereof.

The present invention also relates to novel enzymes which can transform known carotenoids into novel or uncommon products. Currently ϵ -carotene (see Figure 2) and γ -carotene are commonly produced only in minor amounts. As described below, an enzyme can be produced which transforms lycopene to γ -carotene and lycopene to ϵ -carotene. With these products in hand, bulk synthesis of other carotenoids derived from them are possible. For example, ϵ -carotene can be hydroxylated to form lactucaxanthin, an isomer of lutein (one ϵ and one β ring) and zeaxanthin (two β rings) where both endgroups are, instead, ϵ rings.

In addition to novel enzymes produced by truncating the 5' region of known enzymes, as discussed above, novel enzymes which can participate in the formation of unusual carotenoids can be formed by replacing portions of one gene with an analogous sequence from a structurally related gene. For example, β -cyclase and ϵ -cyclase are structurally related (see Figure 13). By replacing a portion of β -lycopene cyclase with the analogous portion of ϵ -cyclase, an enzyme which produces γ -carotene will be produced (one β endgroup). Further, by replacing a portion of the lycopene ϵ -cyclase with the analogous portion of β -cyclase, an enzyme which produces ϵ -carotene will be produced (with some exceptions, such as the lettuce ϵ -cyclase, plant ϵ -cyclases normally produce a compound with one ϵ -endgroup, δ -carotene). Similarly, β -hydroxylase could be modified to produce enzymes of novel function by creation of hybrids with ϵ -hydroxylase.

Host systems according to the present invention can comprise any organism that already produces carotenoids or which has been genetically modified to produce carotenoids.

The IPP isomerase genes are more broadly applicable for enhancing production of any product dependent on DMAPP and/or IPP as a precursor.

5

10

25

30

Organisms which already produce carotenoids include plants, algae, some yeasts, fungi and cyanobacteria and other photosynthetic bacteria. Transformation of these hosts with vectors according to the present invention can be done using standard techniques such as those described in Misawa et al., (1990) supra; Hundle et al., (1993) supra; Hundle et al., (1991) supra; Misawa et al., (1991) supra; Sandmann et al., supra; and Schnurr et al., supra.

Transgenic organisms can be constructed which include the nucleic acid sequences of the present invention (Bird et al, 1991; Bramley et al, 1992; Misawa et al, 1994a; Misawa et al, 1994b; Cunningham et al, 1993). The incorporation of these sequences can allow the controlling of carotenoid biosynthesis, content, or composition in the host cell. These transgenic systems can be constructed to incorporate sequences which allow for the overexpression of the nucleic acids of the present invention. Transgenic systems can also be constructed containing antisense expression of the nucleic acid sequences of the present invention. Such antisense expression would result in the accumulation of the substrates of the substrates of the enzyme encoded by the sense strand.

A method for screening for eukaryotic genes which encode enzymes involved in carotenoid biosynthesis comprises transforming a prokaryotic host with a nucleic acid which may contain a eukaryotic or prokaryotic carotenoid biosynthetic gene; culturing said transformed host to obtain colonies; and screening for colonies exhibiting a different color than colonies of the untransformed host.

Suitable hosts include *E. coli*, cyanobacteria such as *Synechococcus* and *Synechocystis*, alga and plant cells. *E. coli* are preferred.

In a preferred embodiment, the above "color complementation" screening protocol can be enhanced by using mutants which are either (1) deficient in at least one carotenoid biosynthetic gene or (2) overexpress at least one carotenoid biosynthetic gene. In either case, such mutants will accumulate carotenoid precursors.

Prokaryotic and eukaryotic DNA or cDNA libraries can be screened in total for the presence of genes of carotenoid biosynthesis, metabolism and degradation. Preferred organisms to be screened include photosynthetic organisms.

10

Half the man and the man

20 13

25

30

PCT/US99/12121

E. coli can be transformed with these eukaryotic cDNA libraries using conventional methods such as those described in Sambrook et al, 1989 and according to protocols described by the vendors of the cloning vectors.

For example, the cDNA libraries in bacteriophage vectors such as lambdaZAP (Stratagene) or lambda ZIPLOX (Gibco BRL) can be excised en masse and used to transform E.coli.

Transformed E. coli can be cultured using conventional techniques. The culture broth preferably contains antibiotics to select and maintain plasmids. Suitable antibiotics include penicillin, ampicillin, chloramphenicol, etc. Culturing is typically conducted at 15-40°C, preferably at room temperature or slightly above (18-28°C), for 12 hours to 7 days.

Cultures are plated and the plates are screened visually for colonies with a different color than the colonies of the host E. coli transformed with the empty plasmid cloning vector. For example, E. coli transformed with the plasmid, pAC-BETA (described below), produce yellow colonies that accumulate β-carotene. After transformation with a cDNA library. colonies which contain a different hue than those formed by E. coli/pAC-BETA would be expected to contain enzymes which modify the structure or accumulation of β -carotene. Similar E. coli strains can be engineered which accumulate earlier products in carotenoid biosynthesis, such as lycopene, y-carotene, etc.

Having generally described this invention, a further understanding can be obtained by reference to certain specific examples which are provided herein for purposes of illustration only and are not intended to be limiting unless otherwise specified.

EXAMPLE

I. Isolation of β -carotene hydroxylase

Plasmid Construction

An 8.6kb BgIII fragment containing the carotenoid biosynthetic genes of Erwinia herbicola was first cloned in the BamHI site of plasmid vector pACYC184 (chloramphenicol resistant), and then a 1.1kb BamHI fragment containing the E. herbicola β-carotene hydroxylase (CrtZ) was deleted. E.coli strains containing the resulting plasmid, pAC-BETA. accumulate ß-carotene and form yellow colonies (Cunningham et al., 1994).

A full length cDNA encoding IPP isomerase of Haematococcus pluvialis (HP04) was first excised with BamHI and KpnI from pBluescript SK-, and then ligated into the

WO 99/63055 PCT/US99/12121

corresponding sites of the pTrcHisA vector with high-level expression from the *trc* promoter (Invitrogen, Inc.). A fragment containing the IPP isomerase and *trc* promoter was subsequently excised with *EcoRV* and *KpnI*, treated with the Klenow fragment of DNA polymerase to produce blunt ends, and ligated in the Klenow-treated *HindIII* site of pAC-BETA. *E.coli* cells transformed with this new plasmid pAC-BETA-04 form orange colonies on LB plates (*vs.* yellow for those containing pAC-BETA) and cultures accumulate substantially more β-carotene (*ca.* two fold) than those that contain pAC-BETA.

Screening of an Arabidopsis cDNA Library

Several λ cDNA expression libraries of *Arabidopsis* were obtained from the *Arabidopsis* Biological Resource Center (Ohio State University, Columbus, OH) (Kieber et al., 1993). The λ cDNA libraries were excised *in vivo* using Stratagene's ExAssist SOLR system to produce a phagemid cDNA library wherein each phagemid contained also a gene conferring resistance to the antibiotic ampicillin.

E.coli strain DH10BZIP was chosen as the host cell for the screening and pigment production, although we have also used TOP10F' and XL1-Blue for this purpose. DH10B cells were transformed with plasmid pAC-BETA-04 and were plated on LB agar plates containing chloramphenicol at 50 µg/ml (from United States Biochemical Corporation). The phagemid Arabidopsis cDNA library was then introduced into DH10B cells already containing pAC-BETA-04. Transformed cells containing both pAC-BETA-04 and Arabidopsis cDNA library phagemids were selected on chloramphenicol plus ampicillin (150 µg/ml) agar plates. Maximum color development occurred after 3 to 7 days incubation at room temperature, and the rare bright yellow colonies were selected from a background of many thousands of orange colonies on each agar plate. Selected colonies were inoculated into 3 ml liquid LB medium containing ampicillin and chloramphenicol, and cultures were incubated at room temperature for 1-2 days, with shaking. Cells were then harvested by centrifugation and extracted with acetone in microfuge tubes. After centrifugation, the pigmented extract was spotted onto silica gel thin-layer chromatography (TLC) plates, and developed with a hexane:ether (1:1, by volume) mobile phases. B-carotene hydroxylaseencoding cDNAs were identified based on the appearance of a yellow pigment that comigrated with zeaxanthin on the TLC plates.

30

25

5

10

that are a subject to the subject to

30

5

10

Subcloning and Sequencing

The plasmid containing the β-carotene hydroxylase cDNA was recovered and analyzed by standard procedures (Sambrook et al., 1989). The *Arabidopsis* β-carotene hydroxylase was sequenced completely on both strands on an automatic sequencer (Applied Biosystems, Model 373A, Version 2.0.1S). The cDNA insert of 0.95kb also was excised and ligated into the a pTrcHis vector. A *BgI*II restriction site within the cDNA was used to remove that portion of the cDNA that encodes the predicted polypeptide N terminal sequence region that is not also found in bacterial β-carotene hydroxylases (Figure 6). A BgIII-XhoI fragment was directionally cloned in BamHI-XhoI digested TrcHis vectors.

Pigment Analysis

A single colony was used to inoculate 50 ml of LB containing ampicillin and chloramphenicol in a 250-ml flask. Cultures were incubated at 28°C for 36 hours with gentle shaking, and then harvested at 5000 rpm in an SS-34 rotor. The cells were washed once with distilled H₂O and resuspended with 0.5 ml of water. The extraction procedures and HPLC were essentially as described previously (Cunningham et al, 1994).

II. Isolation and biochemical analysis of an Arabidopsis lycopene ϵ -cyclase

Plasmid Construction

Construction of plasmids pAC-LYC, pAC-NEUR, and pAC-ZETA is described in Cunningham et al., (1994). In brief, the appropriate carotenoid biosynthetic genes from *Erwinia herbicola*, *Rhodobacter capsulatus*, and *Synechococcus* sp. strain PCC7942 were cloned in the plasmid vector pACYC184 (New England BioLabs, Beverly, MA). Cultures of *E. coli* containing the plasmids pAC-ZETA, pAC-NEUR, and pAC-LYC, accumulate ζ-carotene, neurosporene, and lycopene, respectively. The plasmid pAC-ZETA was constructed as follows: an 8.6-kb BglII fragment containing the carotenoid biosynthetic genes of *E. herbicola* (GenBank M87280; Hundle et al., 1991) was obtained after partial digestion of plasmid pPL376 (Perry et al., 1986; Tuveson et al., 1986) and cloned in the BamHI site of pACYC184 to give the plasmid pAC-EHER. Deletion of adjacent 0.8- and 1.1-kb BamHI-BamHI fragments (deletion Z in Cunningham et al., 1994), and of a 1.1 kB SalI-SalI fragment (deletion X) served to remove most of the coding regions for the *E. herbicola* β-carotene hydroxylase (crtZ gene) and zeaxanthin glucosyltransferase (crtX gene), respectively. The

25

10

5

resulting plasmid, pAC-BETA, retains functional genes for geranylgeranyl pyrophosphate synthase (crtE), phytoene synthase (crtB), phytoene desaturase (crtI), and lycopene cyclase (crtY). Cells of *E. coli* containing this plasmid form yellow colonies and accumulate β -carotene. A plasmid containing both the lycopene ϵ - and β -cyclase cDNAs of *A. thaliana* was constructed by excising the ϵ -cyclase in clone y2 as a PvuI-PvuII fragment and ligating this piece in the SnaBI site of a plasmid (pSPORT 1 from GIBCO-BRL) that already contained the β -cyclase (Cunningham et al., 1996).

Organisms and Growth Conditions

E. coli strains TOP10 and TOP10 F' (obtained from Invitrogen Corporation, San Diego, CA) and XL1-Blue (Stratagene) were grown in Luria-Bertani (LB) medium (Sambrook et al., 1989) at 37°C in darkness on a platform shaker at 225 cycles per min. Media components were from Difco (yeast extract and tryptone) or Sigma (NaCl). Ampicillin at 150 μg/mL and/or chloramphenicol at 50 μg/mL (both from United States Biochemical Corporation) were used, as appropriate, for selection and maintenance of plasmids.

Mass Excision and Color Complementation Screening of an A. thaliana cDNA Library

A size-fractionated 1-2 kB cDNA library of *A. thaliana* in lambda ZAPII (Kieber et al., 1993) was obtained from the Arabidopsis Biological Resource Center at The Ohio State University (stock number CD4-14). Other size fractionated libraries were also obtained (stock numbers CD4-13, CD4-15, and CD4-16). An aliquot of each library was treated to cause a mass excision of the cDNAs and thereby produce a phagemid library according to the instructions provided by the supplier of the cloning vector (Stratagene; *E. coli* strain XL1-Blue and the helper phage R408 were used). The titre of the excised phagemid was determined and the library was introduced into a lycopene-accumulating strain of *E. coli* TOP10 F' (this strain contained the plasmid pAC-LYC) by incubation of the phagemid with the *E. coli* cells for 15 min at 37°C. Cells had been grown overnight at 30°C in LB medium supplemented with 2% (w/v) maltose and 10 mM MgSO₄ (final concentration), and harvested in 1.5 ml microfuge tubes at a setting of 3 on an Eppendorf microfuge (5415C) for 10 min. The pellets were resuspended in 10 mM MgSO₄ to a volume equal to one-half that of the

30

5

10

initial culture volume. Transformants were spread on large (150 mm diameter) LB agar petri plates containing antibiotics to provide for selection of cDNA clones (ampicillin) and maintenance of pAC-LYC (chloramphenicol). Approximately 10,000 colony forming units were spread on each plate. Petri plates were incubated at 37.C for 16 hr and then at room temperature for 2 to 7 days to allow maximum color development. Plates were screened visually with the aid of an illuminated 3x magnifier and a low power stage-dissecting microscope for the rare, pale pinkish-yellow to deep-yellow colonies that could be observed in the background of pink colonies. A colony color of yellow or pinkish-yellow was taken as presumptive evidence of a cyclization activity. These yellow colonies were collected with sterile toothpicks and used to inoculate 3ml of LB medium in culture tubes with overnight growth at 37°C and shaking at 225 cycles/min. Cultures were split into two aliquots in microfuge tubes and harvested by centrifugation at a setting of 5 in an Eppendorf 5415C microfuge. After discarding the liquid, one pellet was frozen for later purification of plasmid DNA. To the second pellet was added 1.5 ml EtOH, and the pellet was resuspended by vortex mixing, and extraction was allowed to proceed in the dark for 15-30 min with occasional remixing. Insoluble materials were pelleted by centrifugation at maximum speed for 10 min in a microfuge. Absorption spectra of the supernatant fluids were recorded from 350-550 nm with a Perkin Elmer lambda six spectrophotometer.

Analysis of isolated clones

Eight of the yellow colonies contained β -carotene indicating that a single gene product catalyzes both cyclizations required to form the two β endgroups of the symmetrical β -carotene from the symmetrical precursor lycopene. One of the yellow colonies contained a pigment with the spectrum characteristic of δ -carotene, a monocyclic carotenoid with a single ϵ endgroup. Unlike the β cyclase, this ϵ -cyclase appears unable to carry out a second cyclization at the other end of the molecule.

The observation that ϵ -cyclase is unable to form two cyclic ϵ -endgroups (e.g. the bicyclic ϵ -carotene) illuminates the mechanism by which plants can coordinate and control the flow of substrate into carotenoids derived from β -carotene versus those derived from α -carotene and also can prevent the formation of carotenoids with two ϵ endgroups.

The availability of the A. thaliana gene encoding the ϵ -cyclase enables the directed manipulation of plant and algal species for modification of carotenoid content and

30

5

10

composition. Through inactivation of the ϵ -cyclase, whether at the gene level by deletion of the gene or by insertional inactivation or by reduction of the amount of enzyme formed (by such as antisense technology), one may increase the formation of β -carotene and other pigments derived from it. Since vitamin A is derived only from carotenoids with β endgroups, an enhancement of the production of β -carotene versus α -carotene may enhance nutritional value of crop plants. Reduction of carotenoids with ϵ -endgroups may also be of value in modifying the color properties of crop plants and specific tissues of these plants. Alternatively, where production of α -carotene, or pigments such as lutein that are derived from α -carotene, is desirable, whether for the color properties, nutritional value or other reason, one may overexpress the ϵ -cyclase or express it in specific tissues. Wherever agronomic value of a crop is related to pigmentation provided by carotenoid pigments the directed manipulation of expression of the ϵ -cyclase gene and/or production of the enzyme may be of commercial value.

The predicted amino acid sequence of the A. thaliana ϵ -cyclase enzyme was determined. A comparison of the amino acid sequences of the β - and ϵ -cyclase enzymes of Arabidopsis thaliana (Fig. 13) as predicted by the DNA sequence of the respective cDNAs (Fig. 4 for the ϵ -cyclase cDNA sequence), indicates that these two enzymes have many regions of sequence similarity, but they are only about 37% identical overall at the amino acid level. The degree of sequence identity at the DNA base level, only about 50%, is sufficiently low such that we and others have been unable to detect this gene by hybridization using the β cyclase as a probe in DNA gel blot experiments.

REFERENCES

Each reference cited in this application and/or listed below is hereby incorporated by reference.

Bird et al, 1991 Biotechnology 9, 635-639.

Bishop et al., (1995) FEBS Lett. 367, 158-162.

Bramley, P.M. (1985) Adv. Lipid Res. 21, 243-279.

Bramley, P.M. (1992) Plant J. 2, 343-349.

Britton, G. (1988). Biosynthesis of carotenoids. In Plant Pigments, T.W. Goodwin, ed. (London: Academic Press), pp. 133-182.

30

5

10

Britton, G. (1979) Z. Naturforsch. Section C Biosci. 34, 979-985.

Britton, G. (1995) UV/Visible spectroscopy. In Carotenoids, Vol. IB: Spectroscopy,

G. Britton, S. Liaaen-Jensen, H.P. Pfander, eds. (Basel: Birkhauser Verlag), pp. 13-62.

Bouvier et al., (1994) Plant J. 6, 45-54.

Cunningham et al., (1985) Photochem. Photobiol. 42: 295-307.

Cunningham et al., (1993) FEBS Lett. 328, 130-138.

Cunningham et al., (1994) Plant Cell 6, 1107-1121.

Cunningham et al., (1996) Plant Cell 8, 1613-1626.

Davies, B.H. (1976). Carotenoids. In Chemistry and Biochemistry of Plant Pigments,

Vol. 2, T.W. Goodwin, ed (New York: Academic Press), pp. 38-165.

Del Sal et al., (1988). Nucl. Acids Res. 16, 9878.

Demmig-Adams & Adams, (1992) Ann. Rev. Plant Physiol. Mol. Biol. 43, 599-626.

Enzell & Back, (1995) Mass spectrometry. In Carotenoids, Vol. IB: Spectroscopy, G.

Britton, S. Liaaen-Jensen, H.P. Pfander, eds. (Basel: Birkhauser Verlag), pp. 261-320.

Frank & Cogdell (1993) Photochemistry and function of carotenoids in photosynthesis. In Carotenoids in Photosynthesis. A. Young and G. Britton, eds. (London: Chapman and Hall). pp. 253-326.

Goodwin, T.W. (1980). The Biochemistry of the Carotenoids. 2nd ed, Vol. 1 (London: Chapman and Hall.

Horvath et al., (1972) Phytochem. 11, 183-187.

Hugueney et al., (1995) Plant J. 8, 417-424.

Hundle et al., (1991) Photochem. Photobiol. 54, 89-93.

Jensen & Jensen, (1971) Methods Enzymol. 23, 586-602.

Kargl & Quackenbush, (1960) Archives Biochem. Biophys. 88, 59-63.

Kargl et al., (1960) Proc. Am. Hort. Soc. 75, 574-578.

Kieber et al., (1993) Cell 72, 427-441.

Kovama, Y. (1991) J. Photochem. Photobiol., B, 9, 265-80.

Krinsky, N.I. (1987) Medical uses of carotenoids. In Carotenoids, N.I. Krinsky, M.M. Mathews-Roth, and R.F. Taylor, eds. (New York: Plenum), pp. 195-206.

Kyte & Doolittle, (1982) J. Mol. Biol. 157, 105-132.

LaRossa & Schloss, (1984) J. Biol. Chem. 259, 8753-8757.

Misawa et al., (1994a) Plant J. 6, 481-489.

30

5

10

Misawa et al., (1994b) J. Biochem, Tokyo, 116, 980-985.

Norris et al., (1995) Plant Cell 7, 2139-2149.

Pecker et al., (1996) Submitted to Plant Mol. Biol.

Perry et al., (1986) J. Bacteriol. 168, 607-612.

Persson & Argos, (1994) J. Mol. Biol. 237, 182-192.

Plumley & Schmidt, (1987) Proc. Nat. Acad. Sci. USA 83, 146-150.

Plumley & Schmidt, (1995) Plant Cell 7, 689-704.

Rossmann et al., (1974) Nature 250, 194-199.

Rock & Zeevaart (1991) Proc. Nat. Acad. Sci. USA 88, 7496-7499.

Rost et al., (1995) Protein Science 4, 521-533.

Sambrook et al., (1989) Molecular Cloning: A Laboratory Manual, 2nd edition (Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press).

Sancar, A. (1994) Biochemistry 33, 2-9.

Sander & Schneider, (1991) Proteins 9, 56-68.

Sandmann, G. (1994) Eur. J. Biochem. 223, 7-24.

Scolnik & Bartley, (1995) Plant Physiol. 108, 1342.

Siefermann-Harms, D. (1987) Physiol. Plant. 69, 561-568.

Spurgeon & Porter, (1980). Biosynthesis of carotenoids. In Biochemistry of Isoprenoid Compounds, J.W. Porter, and S.L. Spurgeon, eds. (New York: Wiley), pp. 1-122.

Tomes, M.L. (1963) Bot. Gaz. 124, 180-185.

Tomes, M.L. (1967) Genetics 56, 227-232.

Tuveson et al., (1986) J. Bacteriol. 170, 4675-4680.

Van Beeumen et al., (1991) J. Biol. Chem. 266, 12921-12931.

Weedon & Moss, (1995) Structure and Nomenclature. In Carotenoids, Vol. IB:

Spectroscopy, G. Britton, S. Liaaen-Jensen, H.P. Pfander, eds. (Basel: Birkhauser Verlag), pp. 27-70.

Wierenga et al., (1986) J. Mol. Biol. 187, 101-107.

Zechmeister, L. (1962) Cis-Trans Isomeric Carotenoids, Vitamins A and Arylpolyenes. Springer-Verlag, Vienna.

Having now fully described the invention, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the invention as set forth herein.

We claim:

- 1. An isolated and/or purified nucleic acid sequence which encodes for a protein having lycopene ϵ -cyclase enzyme activity and has an amino acid sequence which is at least 85% identical to one of SEQ ID NOS: 23 or 25-27.
- 5 2. The nucleic acid sequence of claim 1, wherein the protein has the amino acid sequence of one of SEQ ID NOS: 23 or 25-27.
 - 3. A vector comprising the nucleic acid sequence of claim 1, wherein the nucleic acid sequence is operably linked to a promoter.
 - 4. A host cell which contains the vector of claim 3.
 - 5. The host cell of claim 4, wherein the host cell is selected from the group consisting of a bacterial cell, an algal cell, a yeast cell and a plant cell.
 - 6. The host cell of claim 4, wherein the host cell is a photosynthetic cell.
 - 7. An isolated and/or purified protein having lycopene ϵ -cyclase enzyme activity and having an amino acid sequence which is at least 85% identical to one of SEQ ID NOS: 23 or 25-27.
 - 8. The protein of claim 7, wherein the protein has the amino acid sequence of one of SEQ ID NOS: 23 or 25-27.

AMENDED CLAIMS

[received by the International Bureau on 15 November 1999 (15.11.99); original claims 1,2,7 and 8 amended; remaining claims unchanged (1 page)]

- 1. An isolated and/or purified nucleic acid sequence which encodes for a protein having lycopene ∈-cyclase enzyme activity and has an amino acid sequence which is at least 85% identical to one of SEQ ID NOS: 23, 25 or 26.
- 5 2. The nucleic acid sequence of claim 1, wherein the protein has the amino acid sequence of one of SEQ ID NOS: 23, 25 or 26.
 - 3. A vector comprising the nucleic acid sequence of claim 1, wherein the nucleic acid sequence is operably linked to a promoter.
 - 4. A host cell which contains the vector of claim 3.
 - 5. The host cell of claim 4, wherein the host cell is selected from the group consisting of a bacterial cell, an algal cell, a yeast cell and a plant cell.
 - 6. The host cell of claim 4, wherein the host cell is a photosynthetic cell.
 - 7. An isolated and/or purified protein having lycopene ϵ -cyclase enzyme activity and having an amino acid sequence which is at least 85% identical to one of SEQ ID NOS: 23, 25 or 26.
 - 8. The protein of claim 7, wherein the protein has the amino acid sequence of one of SEQ ID NOS: 23, 25 or 26.

1/45

FIG.1

2/45

desolurase desolurase
$$\frac{1}{2}$$
 $\frac{1}{4}$ $\frac{1}{6}$ $\frac{1}{8}$ $\frac{1}{4}$ $\frac{1}{6}$ $\frac{1}{8}$ $\frac{1}{6}$ $\frac{1}{4}$ $\frac{1}{6}$ $\frac{1}{8}$ $\frac{1}{6}$ $\frac{1}{8}$ $\frac{$

3/45

$$\frac{2}{3} + \frac{6}{4}$$

$$\psi \text{ endgroup}$$

$$\beta \text{ cyclase}$$

$$\frac{2}{4} + \frac{6}{5}$$

$$\frac{2}{4} + \frac$$

4/45

09/701395

FIG.4A

F1G.4B

F1G. 4

FIG. IIA

FIG.11B

FIG. 11

FIG. 13A

FIG. 13B

FIG. 13

FIG.14A

FIG. 14B

FIG. 14

FIG. 22A

FIG. 22B

FIG. 22

FIG. 4A

Arabidopsis thaliana epsilon cyclase:

				а	caa	laag	i gaa	ata	att	ag	att	cct	ctti	tctg	gct	tgc	tata	acct	tga	.ca	48
	gaac	aac	ata	laca	atç	gto	jtaa	igto	ttc	tc	gct	gta	ttc	gaaa	att	atti	tgga	agga	igga	ac	108
1	atgg M	jagt E	gto C	jtte V	igg99	icta A	igga R	att N	tcg F	jca A	gca: A	atg M	gcgg	gtti V	tca S	aca [*] T	ttto F	ccgt P	cat S	.gg W	168
21	agti S	igto C	cgaa R	agga R	aat K	tto F	ccag P	gtgg V	octa V	aag K	aga R	tac Y	agc S	tata Y	agg R	aata N	atto I	cgc1 R	tcg F	gt G	228
41	ttg L	tgta C	agto S	gtca V	agag R	gcta A	agc(ggcg G	g G	gga G	agt S	tco S	ggt. G	agto S	gag K	agt: S	tgt:	gtaq V	gcgg A	rtg V	288
61	agag	gaa S	gat D	tto: F	octo A	gaci D	gaaq E	gaaq E	gati D	ttt F	gcg V	aaa E	gct: A	G ggci	ggt: G	tct S	gaga R	atto I	ctat L	tt F	348
81	gtt V	caa Q	atg M	cago Q	caga Q	aaca M	aaa K	gata D	atgo M	gat D	gaa S	.cag Q	tct S	aag K	ctt L	gtt V	gat D	aagi K	tgc L	ct P	408
.01	cct P	ata I	tca S	act I	ggt G	gat D	ggt G	gct [.] A	ttg L	gat D	cat K	.gtg V	gtt V	act I	ggc G	tgt C	ggt. G	cct(P	gcts A	ıgt G	468
.21	tta L	igcc A	ttg L	gct A	gca A	gaa K	tca S	gct A	aag K	ctt L	gga G	itta L	aaa K	gtt V	gga G	ctc L	att I	ggt: G	ccag P	jat D	528
141	ctt L	cct P	ttt F	act T	aac M	aat M	tac Y	ggt G	gtt V	tgg M	gaa K	agat D	.gaa K	ttc F	aat N	gat D	ctt L	999 G	ctgo L	zaa G	588
161	aaa K	atgt C	tatt I	gag K	icat K	gtt V	tgg W	jaga R	.gag S	act T	att I	gcg V	icac Y	ctg L	gat D	gat D	gac D	aag K	ccta P	att I	648
181	aco T	cati I	tgga G	ccgt R	gct A	tat Y	:gga G	aga R	gtt V	agt S	cga R	acgt R	ttg L	ctc L	cat X	.gag E	gag E	ctt	ttga L	agg R	708
201	agg R	gtg C	tgto V	cgaç K	jtca S	aggt G	gto V	tcg S	rtac Y	ctt L	ago S	ctco	jaaa K	igtt V	gac D	ago S	ata I	aca T	gaag E	gct A	768
221	tg . S	tga D	tgg G	cct1 L	taga X	acti L	tgt1 V	tgct A	tgt C	gac D	gad D	caat M	taac M	gto V	att I	CCC P	tgc C	agg X	ctto L	gcc A	828
241	ac T	tgt V	tgc A	ttc S	tgg: G	agca A	agci A	ttcg S	999a G	aaag K	cto	ctt L	gcaa Q	atac Y	gaa X	igtt V	.ggt G	.gga G	ccta P	aga R	888
	c rt	· / + /-	rtac	aca.	aac	tac	ata	COCK	ata	ndac	at:	t.aac		ogaa	aaat	aat	.cc	itat	gate	cca	948

F1G. 4B

261	V C	٧	Q	Т	A	Y	G	٧	Χ	٧	X	٧	χ	N	S	P	Υ	D	P	
281	gatca D Q	aato M	ggtt V	ttc P	atg M	gat D	tac Y	caga R	agat D	tat Y	act T	aac M	gag X	jaaa X	gtt V	.cgg R	jago S	tta L	agaa X	1008
301	gctga A K	gta Y	tcca p	acg T	rttt F	ctg L	tac Y	gcc A	atg M	cct P	atg M	aca T	aag K	rtca S	aga R	icto L	tto F	ttc F	gag K	1068
321	gagac K T	atg C	tttg L	99CC A	tca S	iaaa K	gat D	gto V	atg M	ccc P	ttt F	gat D	ttg L	icta L	iaaa K	acg T	jaag K	gcto L	atg M	1128
341	ttaag I F	att V	agad G	caca G	cto S	:gga L	att P	cga N	aatt T	cta X	aag Q	act K	tac N	:gaa L	igag A	gag F	rtgg G	rtcc A	tat A	1188
361	atccc I F	agt V	tggt G	.ggt G	tcc S	ttg L	KCZ P	aaa M	cacc T	gaa X	caa Q	aag K	aat N	ctc L	gcc A	ttt F	.ggt G	gct A	gcc A	1248
381	gctac A S	icat M	ggta V	acat M	CCC P	:gca A	aca T	aggo G	tat Y	tca S	gtt V	gtg V	aga R	tct S	ttg L	tct S	gaa X	igct A	cca P	1308
401	aaaca K	atgc / A	atca S	agto V	ato I	gca A	gag K	gata I	acta L	aga R	gaa E	gag E	act T	acc T	aaa K	cag Q	att I	aac N	agt S	1368
421	aatat M]	itto I S	aaga R	acaa Q	agct A	tag W	gat D	tact T	tta L	tgg W	cca P	.cca P	gaa E	agg R	iaaa X	aga R	.cag Q	jaga R	gca A	1428
441	ttct! F f	ttct L	ctt	tggt G	ctt L	igca A	cto L	caga I	agtt V	caa Q	ttc F	gat D	acc T	gaa X	.ggc G	att I	aga R	agc S	ttc F	1488
461	ttcc F	gtac R T	ttt	ctto F	ccgc R	ctt L	cca p	aaaa K	atgg W	atg M	tgg W	caa Q	.ggg G	ttt F	cta L	gga G	tca S	aca T	tta L	1548
481	acato	cago S G	iaga D	tcto L	gt1 V	tctc L	tt! F	tgci A	ttta L	tac Y	atg M	rttc P	gtc V	att I	tca S	cca P	aac M	aat M	ttg L	1608
501	agaa R I	aagg K	rtct	catt I	taat N	cat W	cto	cato I	ctct S	gat D	.cca P	acc T	gga G	gca A	acc T	atg M	ata I	aaa K	acc T	1668
521	tato	tcaa L k	_	atga	atti	tact	ta	cca	actc	tta	.ggt	ttg	tgt	ata	tat	atg	ccg	att	tat	1728
<i>ح</i> يد	ctg <u>a</u>			atca	aaa	gaat	gg	tate	gtgg	gtt	act	agg	jaag	rttg	gaa	aca	aac	acq	tat	1788
	agaa		_											_	_					1848
	ccgt	ggct	agt	g																1868

F1G. 5

gctctttctc ctcctctt accgatttcc gactccgcct cccgaaatcc 1 ttatccggat tctctccgtc tcttcgattt aaacgctttt ctgtctgtta 51 cgtcgtcgaa gaacggagac agaattctcc gattgagaac gatgagagac 101 cggagagcac gagctccaca aacgctatag acgctgagta tctggcgttg 151 cgtttggcgg agaaattgga gaggaagaaa tcggagaggt ccacttatct 201 aatcgctgct atgttgtcga gctttggtat cacttctatg gctgttatgg 251 ctgtttacta cagattctct tggcaaatgg agggaggtga gatctcaatg 301 ttggaaatgt ttggtacatt tgctctctct gttggtgctg ctgttggtat 351 ggaattctgg gcaagatggg ctcatagagc tctgtggcac gcttctctat 401 ggaatatgca tgagtcacat cacaaaccaa gagaaggacc gtttgagcta 451 aacgatgttt ttgctatagt gaacgctggt ccagcgattg gtctcctctc 501 ttatggattc ttcaataaag gactcgttcc tggtctctgc tttggcgccg 551 ggttaggcat aacggtgttt ggaatcgcct acatgtttgt ccacgatggt 601 ctcgtgcaca agcgtttccc tgtaggtccc atcgccgacg tcccttacct 651 ccgaaaggtc gccgccgctc accagctaca tcacacagac aagttcaatg 701 qtqtaccata tqqactgttt cttggaccca aggaattgga agaagttgga 751 ggaaatgaag agttagataa ggagattagt cggagaatca aatcatacaa 801 aaaggcctcg ggctccgggt cgagttcgag ttcttgactt taaacaagtt 851 ttaaatccca aattctttt ttgtcttctg tcattatgat catcttaaga 901 951 caatct

	SFSS SSTDFRLRLP KSLSGFSPSL RFKRFSVCYV VEERRQNSPI ENDERPESIS SIIVALDAETL 144	ALRLAEKLER KKSERSTYLI AAMLSSFGIT SWAWAVYYR FSWQMEGGEI SMLEMFGIFA LSVGAAVGME FWARWAHRAL MTOFL IVVATVLWE LTAYSVHRWI MINFL IVVATVLWE LTAYSVHRWI ML.NSL IVILSVIAME GIAAFTHRYI MLWIWNAL IVIVTVIGME VIAALAHKYI MLWIWNAL IVIVTVIGME VIAALAHKYI	Predicted TM helix	WHASL.WNMH ESHHKPREGP FELNDVFAIV NAGPAIGLLS YGFFNKGLVP GLCFGAGLGI TVFGIAYMFV HDGLVHKRFP MFGPLGWGMH KSHHEEHDHA LEKNDLYGVV FAVLATILFT VGAYWMPVLW WIALGM TVYGLIYFIL HDGLVHQRWP MFGPLGWGMH KSHHEEHDHA LEKNDLYGLV FAVIATVLFT VGWIWAPVLW WIALGM TVYGLIYFVL HDGLVHQRWP MG.WGWRMH ESHHTPRKGV FKLNDLFAVV FAGVAIALIA VGTAGWWPLQ WIGCGM TVYGLLYFLV HDGLVHQRWP MFG.WGWGWH LSHHEPRKGA FEVNDLYAVV FAGVAIALIY LGSTGMWPLQ WIGAGM TAYGLLYFLV HDGLVHQRWP HLSHHEPRKGA FEVNDLYAVV FAALSILLIY LGSTGMWPLQ WIGAGM TAYGLLYFNV HDGLVHQRWP -H1-WH -SHH-pr-g- fE-NDa-V -Aai-LGglG- TV-GYV HDGLVH-R-P	Predicted TM helix Predicted TM helix	301 VGPIADVPYL RKVAAAHOLH HT. DKFNGV PYGLFLGPKE LEEVGGNEEL DKEISRRIKS YKKASGSGSS SSS* FRYIPRRGYF RRLYQAHRLH HAVEGROHCV SFGFIYAPP. VDKLKQDLKR SGVLRPQDER PS*
F1G. 6	A.thal.	A.thal. A Alical. A.aurant. E.herb. E.ured. Consensus		A.thal. M.aurant. N. E.herb. E.ured. N. Consensus		A.thal. Alical. A.aurant. E.herb. E.ured. Consensus

9/45

FIG. 7

ccacqqqtcc qcctccccqt ttttttccqa tccqatctcc qqtqccqaqq actcagetgt ttgttegege ttteteagee gtcaccatga eegattetaa 51 cgatgctgga atggatgctg ttcagagacg actcatgttt gaagacgaat 101 gcattetegt tgatgaaaat aategtgtgg tgggacatga cactaagtat 201 aactgtcatc tgatggaaaa gattgaagct gagaatttac ttcacagagc 251 tttcagtgtg tttttattca actccaagta tgagttgctt ctccagcaac 301 ggtcaaaaac aaaggttact ttcccacttg tgtggacaaa cacttqttgc agccatcoto tttacogtga atcogagott attgaagaga atgtgcttgg 351 401 tgtaagaaat googoacaaa ggaagotttt ogatgagoto ggtattgtag 451 cagaagatgt accagtogat gagttcacto cottgggacg catgotttac 501 aaggcacctt ctgatgggaa atggggagag cacgaagttg actatctact 551 cttcatcgtg cgggatgtga agcttcaacc aaacccagat gaagtggctg 601 agatcaagta cgtgagcagg gaagagctta aggagctggt gaagaaagca 651 gatgetggeg atgaagetgt gaaactatet ceatggttea gattggtggt 701 ggataatttc ttgatgaagt ggtgggatca tgttgagaaa ggaactatca 751 ctgaagctgc agacatgaaa accattcaca agctctgaac tttccataag 801 tittggatct toccottocc ataataaaat taagagatga gacttttatt 851 gattacagac aaaactggca acaaaatcta ttcctaggat ttttttttcc 901 tttttattta cttttgattc atctctagtt tagttttcat cttaaaaaaa 951 aaaa

FIG. 8

caccaatgto tgtttottot ttatttaato toocattgat togootoaga 51 totatagate titagiate tittatict titaGATTTG CCCATCGTCC 101 TETGTCATCG ATTTCACCGA GAAAGTTACC GAATTTTCGT GCTTTCTCTG GTACCGCTAT GACAGATACT AAAGATGCTG GTATGGATGC TGTTCAGAGA 151 CGTCTCATGT TTGAGGATGA ATGCATTCTT GTTGATGAAA CTGATCGTGT 201 TGTGGGGCAT GTCAGCAAGT ATAATTGTCA TCTGATGGAA AATATTGAAG 251 301 CCAAGAATTT GCTGCACAGG GCTTTTAGTG TATTTTTATT CAACTCGAAG 351 TATGAGTTGC TTCTCCAGCA AAGGTCAAAC ACAAAGGTTA CGTTCCCTCT AGTGTGGACT AACACTTGTT GCAGCCATCC TCTTTACCGT GAATCAGAGC 401 TTATCCAGGA CAATGCACTA GGTGTGAGGA ATGCTGCACA AAGAAAGCTT 451 CTCGATGAGC TTGGTATTGT AGCTGAAGAT GTACCAGTCG ATGAGTTCAC 501 TCCCTTGGGA CGTATGCTGT ACAAGGCTCC TTCTGATGGC AAATGGGGAG AGCATGAACT TGATTACTTG CTCTTCATCG TGCGAGACGT GAAGGTTCAA 601 CCAAACCCAG ATGAAGTAGC TGAGATCAAG TATGTGAGCC GGGAAGAGCT GAAGGAGCTG GTGAAGAAAG CAGATGCAGG TGAGGAAGGT TTGAAACTGT CACCATGGTT CAGATTGGTG GTGGACAATT TCTTGATGAA GTGGTGGGAT 751 CATGTTGAGA AAGGAACTTT GGTTGAAGCT ATAGACATGA AAACCATCCA CAAACTCTGA ACATCTTTTT TTAAAGTTTT TAAATCAATC AACTTTCTCT TCATCATTTT TATCTTTTCG ATGATAATAA TTTGGGATAT GTGAGACACT TACAAAACTT CCAAGCACCT CAGGCAATAA TAAAGTTTGC GGCCGC

FIG. 9

1	CTCGGTAGCT	GGCCACAATC	GCTATTTGGA	ACCTGGCCCG	GCGGCAGTCC
51	GATGCCGCGA	TGCTTCGTTC	GTTGCTCAGA	GGCCTCACGC	ATATCCCCCG
101	CGTGAACTCC	GCCCAGCAGC	CCAGCTGTGC	ACACGCGCGA	CTCCAGTTTA
151	AGCTCAGGAG	CATGCAGATG	ACGCTCATGC	AGCCCAGCAT	CTCAGCCAAT
201	CTGTCGCGCG	CCGAGGACCG	CACAGACCAC	ATGAGGGGTG	CAAGCACCTG
251	GGCAGGCGGG	CAGTCGCAGG	ATGAGCTGAT	GCTGAAGGAC	GAGTGCATCT
301	TGGTGGATGT	TGAGGACAAC	ATCACAGGCC	ATGCCAGCAA	GCTGGAGTGT
351	CACAAGTTCC	TACCACATCA	GCCTGCAGGC	CICCICCYCC	GGGCCTTCTC
401	TGTGTTCCTG	TTTGACGATC	AGGGGCGACT	GCTGCTGCAA	CAGCGTGCAC
451	GCTCAAAAAT	CACCTTCCCA	AGTGTGTGGA	CGAACACCTG	CTGCAGCCAC
501	CCTTTACATO	GGCAGACCCC	AGATGAGGTG	GACCAACTAA	GCCAGGTGGC
551	CGACGGAACA	GTACCTGGCG	CAAAGGCTGC	TGCCATCCGC	AAGTTGGAGC
601		GATACCAGCG			
651		TGCACTACTG			
701		TGGGGCGAGC			
751		c crreececc			
801		G AGGAGCTGCG			
851		G TGGTTTCGCA			
901		T GGACGCGGCC			
951		C ACATCAACGA			
1001		T GGGGTGGAA1			
1051		G AACCTGCAGT			
1101		T ACCGTCACT	r TTTGTCGCGI	ATACTGAACI	· CCAAGAGGTC
1151	*******	LALLA A			

FIG. 10

1	CTCGGTAGCT	GGCCACAATC	GCTATTTGGA	ACCTGGCCCG	GCGGCAGTCC
51	GATGCCGCGA	TGCTTCGTTC	GTTGCTCAGA	GGCCTCACGC	ATATCCCGCG
101	CGTGAACTCC	GCCCAGCAGC	CCAGCTGTGC	ACACGCGCGA	CTCCAGTTTA
151	AGCTCAGGAG	CATGCAGCTG	CTTTCCGAGG	ACCGCACAGA	CCACATGAGG
201	GGTGCAAGCA	CCTGGGCAGG	CGGGCAGTCG	CAGGATGAGC	TGATGCTGAA
251	GGACGAGTGC	ATCTTGGTAG	ATGTTGAGGA	CAACATCACA	GGCCATGCCA
301	GCAAGCTGGA	GTGTCACAAG	TTCCTACCAC	ATCAGCCTGC	AGGCCTGCTG
351	CACCGGGCCT	TCTCTGTGTT	CCTGTTTGAC	GATCAGGGGC	GACTGCTGCT
401	GCAACAGCGT	GCACGCTCAA	AAATCACCTT	CCCAAGTGTG	TGGACGAACA
451	CCTGCTGCAG	CCACCCTTTA	CATGGGCAGA	CCCCAGATGA	GGTGGACCAA
501	CTAAGCCAGG	TGGCCGACGG	AACAGTACCT	GGCGCAAAGG	CTGCTGCCAT
551	CCGCAAGTTG	GAGCACGAGC	TGGGGATACC	AGCGCACCAG	CTGCCGGCAA
601	GCGCGTTTCG	CTTCCTCACG	CGTTTGCACT	ACTGTGCCGC	GGACGTGCAG
651		CACAATCAGC			
701		CGGGCCAACG			
751		GTACGTGACG			
801		C TTCAATGGTC			
851		T TGGTGGGCTC			
901		G GGGAACGGT			
951		G TGAAGACAC			
1001		C TTTTTCTGAG			
1051		C ATCGTCTCT			TAGCTAGAGT
1101	CACTGATGA	A TOOTTTACA	A CTTTCAAAA	A AAAAA	

13/45

FIG. 11A

1 HPO4 MLRSLLRGLT HPO5 MLRSLLRGLT ATDP7 MSVSSLFNLP C.brew. MS.SSMLNFT ATOP5 S.cerevMTADNNSM	HIPRVNSAQQ HIPRVNSAQQ .LIRLRSLA. .ASRIVSLPL .TGPPPRFFP PHGAVSSYAK	PSCAHARLQF PSCAHARLQF LSSSFSSFRF LSSPPSRVHL IRSPVPRTQL LVQNQTPEDI	KLRSMOMTLM KLRSMOLL AHRPLSSIS. PLCFFSPISL FVRAFSAV LEEFPEIIPL	QPSISANLSR PRKLPNFRAF TQRFSAKLTF QQRPNTR
SSQATT.MGE T.MTD		DELMLKDECI DELMLKDECI RRLMFEDECI RRLMFEDECI RRLMFEDECI QIKLMNENCI	LVDVEDNITG LVDVEDNITG LVDETDRVVG LVDENDKVVG LVDENNRVVG VLDWDDNAIG	100 HASKLECHKF HASKLECHKF HVSKYNCHLM HESKYNCHLM HDTKYNCHLM AGTKKVCHLM
101 LPHOPAGLLH LPHOPAGLLH ENIEAKNLLH ENIESENLLH EKIEAENLLH ENIE.KGLLH	RAFSVFLFDD RAFSVFLFNS RAFSVFLFNS RAFSVFLFNS	KYELLLQQRS	RSKITFPSVW RSKITFPSVW NTKVTFPLVW ATKVTFPLVW KTKVTFPLVW TEKITFPDLW	150 TNTCCSHPLH TNTCCSHPLH TNTCCSHPLY TNTCCSHPLY TNTCCSHPLY TNTCCSHPLC
151 GQTPDEVDQL GQTPDEVDQL RE RE IDDELGL	SQVADGTVPG SELIQDNALG SELIDENCLG SELIEENVLG	AKAAAIRKLE VRNAAQRKLL VRNAAQRKLL	DELGIVAEDV	200 PA.SAFRFLT PA.SAFRFLT PV.DEFTPLG PV.DQFIPLS PV.DEFTPLG KTRGKFHFLN
201 RLHYCAADVO RLHYCAADVO RMLY RILY RMLY	PAATQSALWG .KAPSDGKWG .KAPSDGKWG .KAPSDGKWG	EHELDYLLFI	RANVTL VRDVKV IRDVNL VRDVKL	OPNPDEVAEI DPNPDEVAEV QPNPDEVAEI

14/45

FIG. IIB

300
RYVTQEELRQ MMQ...PDN GLQWSPWFRI IAARFLERWW ADLDAALNTD
RYVTQEELRQ MMQ...PDN GLQWSPWFRI IAARFLERWW ADLDAALNTD
KYVSREELKE LVKKADAGEE GLKLSPWFRL VVDNFLMKWW DHVEKGTLVE
KYMNRDDLKE LLRKADAEEE GVKLSPWFRL VVDNFLFKWW DHVEKGSLKD
KYVSREELKE LVKKADAGDE AVKLSPWFRL VVDNFLMKWW DHVEKGTITE
KWVSPNDLKT MF....ADP SYKFTPWFKI ICENYLFNWW EQLDDLSEVE

301
KHEDWGTVHH INEA*
KHEDWGTVHH INEA*
A.IDMKTIHK L*
A.ADMKTIHK L*
A.ADMKTIHK L*
A.ADMKTIHK L*
NDRQ...IHR ML*


F I G. 12

ccaaaaacaa ctcaaatctc ctccgtcgct cttactccgc catgggtgac 51 gactooggea tggatgotgt toagogaogt otcatgtttg acqatgaatg 101 cattttggtg gatgagtgtg acaatgtggt gggacatgat accaaataca 151 attytcactt gatggagaag attgaaacag gtaaaatgct gcacagagca 201 ttcagcgttt ttctattcaa ttcaaaatac gagttacttc ttcagcaacq 251 stetscaace aagstsacat tteetttast atssaccaae acctsttsca 301 gecatecact ctacagagaa teegagettg tteeegaaac geetgagaga 351 401 451 501 551 601 651 XXXXXXXXX XXXXXXXXX XXXXXXXXX tcatgtgcaa aagggtacac 701 tcactgaatg caatttgata tgaaaaccat acacaagctg atatagaaac 751 acacceteaa eegaaaagea ageetaataa ttegggttgg gtegggteta ccatcaattg tttttttctt ttaacaactt ttaatctcta tttgagcatg 801 ttgattottg tottttgtgt gtaagatttt gggtttcgtt tcagttgtaa 851 901 taatgaacca ttgatggttt gcaatttcaa gttcctatcg acatgtagtg 951 atctaaaaaa

F1G. 13 A

16/45

70	IFA AMAVSTFPSW SCRRKFPVVK RYSYRNÍRFG LCSVRASGGG SSGSESCVAV REDFADEXDF	Cyanobacterial enzyme begins — 140 NKSSALLa LVPETKKKNL DFELPmYDp. S.Kg-VV DLAVVGGGPA GLAVAQQVSE AGLSVCSIDp VKSSALLa LVPETKKKNL DFELPmYDp. S.Kg-VV DLAVVGGGPA GLALAAESAK LGLKVGLIGP VKAGGSEIL. FVQMQQNKDM DEQSKLVDKL ppisiGDGAL DHVVIGCGPA GLALAAESAK LGL-VI-P VKSLVGL-VI-P	Dinucleotide-binding signature	141 -PKLIWPNN YGVWVDEFEA MDLLDCLDaT WSGa-VYiDd -t-KDL-RPY GRVNRKQLKS KMMOKCI-NG DLPFTNN YGVWEDEFND LGLQKCIEHV WRETIVYLDD DKPITIGRAY GRVSRRLLHE ELLRRCVESG PNN YGVW-DEFLC WVY-DDR-Y GRV-RLCG	780	211 VKFHgaKVik ViHE.E-kSm liCnDG-tIQ AtVVLDATGF SR. LVQYDK PYnPGY.QVA YGILAEVEEH VSYLSSKVDS ITKASDGLRL VACDDNNVIP CRLATVASGA ASGKLLQYEV GGPRVCVQTA YGVEVENS VKVQ-A YGGA-GL-QYQ-A YGGV
	1 MECVGAŘNFA ÁMÁVSTFPSW SCRRKFPVVK RYS	Cyanobacterial enzyme begins——SSALLa LVPETKKKNL DFELPmYDp. S.Kg-VGGSEIL. FVQMQQNKDM DEQSKLVDKL ppisiGDG	Possible subunit interaction domain	PKLIWPNN YGVWVDEFEA MDLLDCLDAT WS P. FTNN YGVWEDEFND LGLQKCIEHV WF PNN YGVW-DEFLC W-	Conserved region #1	(FHgaKVik ViHE.E-kSm liCnDG-tIQ A SYLSSKVDS ITKASDGLRL VACDDNNVIP C
	Plant beta MEC A.t.epsilon MEC Consensus		Possi	Plant beta A.t.epsilon Dl Consensus		Plant beta VI A.t.epsilon VI Consensus V



18/45

FIG. 14A

Adonis palaestina E-cyclase cDNA #5 Length: 1898

aaaggagtgt totattaatg ttactgtcgc attottgcaa cacttatatt 51 caaactccat tttcttcttt tctcttcaaa acaacaaact aatgtgagca 101 gagtatctgg ctatggaact acttggtgtt cgcaacctca tctcttcttg 151 ccctgtgtgg acttttggaa caaqaaacct tagtagttca aaactagctt 201 ataacataca tcgatatggt tcttcttgta gagtagattt tcaagtgaga 251 gctgatggtg gaagcgggag tagaagttct gttgcttata aagagggttt 301 tqtqqatqaa qaggatttta tcaaagctqq tqqttctqaq cttttqtttq 351 tccaaatgca gcaaacaaag tctatggaga aacaggccaa gctcgccgat 401 aagttgccac caataccttt tggagaatcc gtgatggact tggttgtaat 451 aggttgtgga cctgctggtc tttcactggc tgcagaagct gctaagctag 501 ggttgaaagt tggccttatt ggtcctgatc ttccttttac aaataattat 551 ggtgtgtggg aagacgagtt caaagatctt ggacttgaac gttgtatcga 601 gcatgcttgg aaggacacca tcgtatatct tgataatgat gctcctgtcc 651 ttattggtcg tgcatatgga cgagttagtc gacatttgct acatgaggag 701 ttgctgaaaa ggtgtgtgga gtcaggtgta tcatatctgg attctaaagt 751 ggaaaggatc actgaagctg gtgatggcca tagccttgta gtttgtgaaa 801 atgagatett tateeettge aggettgeta etgttgeate tggageaget 851 tcagggaaac ttttggagta tgaagtaggt ggccctcgtg tttgtgtcca 901 aaccgcttat ggggtggagg ttgaggtgga gaacaatcca tacgatccca 951 acttaatggt attcatggac tacagagact atatgcaaca gaaattacag 1001 tgctcggaag aagaatatcc aacatttctC tatgtcatgc ccatgtcgcc 1051 aacaagactt ttttttgagg aaacctgttt ggcctcaaaa gatgccatgc 1101 cattcgatct actgaagaga aaactgatgt cacgattgaa gactctgggt 1151 atccaagtta caaaagttta tgaagaggaa tggtcatata ttcctgttgg 1201 tggttcttta ccaaacacag agcaaaagaa cctagcattt ggtqctqcaq 1251 caagcatggt gcatccagca acaggctatt cggttgtacg gtcactgtca 1301 gaagetecaa aatatgette tgtaattgea aagattttga aqeaagataa 1351 ctctgcgtat gtggtttctg gacaaagtag tgcagtaaac atttcaatgc 1401 aagcatggag cagtctttgg ccaaaggagc gaaaacgtca aagagcatTc 1451 tttcttttTg gattagagct tattgtgcag ctagatattg aagcaaccag 1501 aacattettt agaacettet teegettgee aacttggatg tggtggggtt 1551 tccttgggtc ttcactatca tctttcgatc tcgtcttgtt ttccatgtac 1601 atgittgitt tggcgccaaa cagcatgagg atgicacttg tgagacattt 1651 gctttcagat ccttctggtg cagttatggt aagagcttac ctcqaaaggt 1701 agtotoatot attattaaac totagtgttt caccaaataa atgaggatoc 1751 ttcqaatgtg tatatgatca tctctatgta tatcctgtac tctaatctca taaagtaaat gccgggtttg atattgttgt gtcaaaccgg ccaatgatat 1801 1851 aaagtaaatt tattgataca aaagtagttt ttttccttaa aaaaaaaa

19/45

FIG. 14B

Adonis palaestina ε-cyclase #5 predicted polypeptide TRANSLATE from: 113 to: 1702 Length: 529 amino acids

MELLGVRNLI SSCPVWTFGT RNLSSSKLAY NIHRYGSSCR VDFQVRADGG SGSRSSVAYK EGFVDEEDFI KAGGSELLFV QMQQTKSMEK QAKLADKLPP 51 IPFGESVMDL VVIGCGPAGL SLAAEAAKLG LKVGLIGPDL PFTNNYGVWE 101 DEFKDLGLER CIEHAWKDTI VYLDNDAPVL IGRAYGRVSR HLLHEELLKR 151 201 CVESGVSYLD SKVERITEAG DGHSLVVCEN EIFIPCRLAT VASGAASGKL LEYEVGGPRV CVQTAYGVEV EVENNPYDPN LMVFMDYRDY MOOKLOCSEE 251 EYPTFLYVMP MSPTRLFFEE TCLASKDAMP FDLLKRKLMS RLKTLGIOVT 301 351 KVYEEEWSYI PVGGSLPNTE QKNLAFGAAA SMVHPATGYS VVRSLSEAPK YASVIAKILK QDNSAYVVSG QSSAVNISMQ AWSSLWPKER KRQRAFFLFG 401 LELIVOLDIE ATRTFFRTFF RLPTWMWGF LGSSLSSFDL VLFSMYMFVL 451 APNSMRMSLV RHLLSDPSGA VMVRAYLER* 501

20/45

FIG. 15A

DNA sequence of potato cDNA (GenBank R27545) obtained from Nicholas J. Provart

potato.seq Length: 1378 August 2, 1996 13:06 Type: N Check: 605 ... tagcggnnnn naggatgagt tcaaagatct tggtcttcaa gcctgcattg 51 aacatgtttg gcgggatacc attgtatatc ttgatgatga tgatcctatt 101 cttattggcc gtgcctatgg aagagttagt cgccatttac tgcacgagga 151 gttactcaaa aggtgtgtgg aggcaggtgt tttgtatcta aactcgaaag 201 tggataggat tgttgaggcc acaaatggcc acagtcttgt agagtqcqaq 251 ggtgatgttg tgattccctg caggtttgtg actgttgcat cgggagcagc 301 ctcggggaaa ttcttgcagt atgagttggg aggtcctaga gtttctgttc 351 aaacagctta tggagtggaa gttgaggtcg ataacaatcc atttgacccq 401 agcctgatgg ttttcatgga ttatagagac tatgtcagac acgacgctca 451 atctttagaa gctaaatatc caacatttct ctatgccatg cccatgtctc 501 caacacgagt ctttttcgag gaaacttgtt tggcttcaaa agatgcaatg 551 ccattcgatc tgttaaagaa aaaattgatg ttacgattga acaccctcgg 601 tgtaagaatt aaagaaattt atgaggagga atggtcttac ataccagttg 651 gaggatettt gecaaataea gaacaaaaaa eaettgeatt tggtgetget 701 gctagcatgg ttcatccagc cacaggttat tcagtcgtca gatcactgtc 751 tgaagctcca aaatgcgcct tcgtgcttgc aaatatatta cgacaaaatc 801 atagcaagaa tatgcttact agttcaagta ccccgagtat ttcaactcaa 851 gcttggaaca ctctttggcc acaagaacga aaacgacaaa gatcgtttt 901 cctatttgga ctggctctga tattgcagct ggatattgag gggataaggt 951 catttttccg cgcgttcttc cgtgtgccaa aatggatgtg gcagggattt 1001 cttggttcaa gtctttcttn agcagacctc atgttatttg ccttctacat 1051 gtttattatt gcaccaaatg acatgagaag aggcttaatc agacatcttt 1101 tatctgatcc tactggtgca acattgataa gaacttatct tacattttag 1151 agtaaattcc tcctacaata gttgttgaan nagaggcctc attacttcag 1201 attcataaca gaaatcgcgg tctctcgagg ccttgtatat aacattttca 1251 ctaggttaat attgcttgaa taagttgcac agtttcagtt tttgtatctg 1301 cttctttttt gtccaagatc atgtattgan ccaatttata tacattgcca 1351 gtatatataa attttataaa aaaaaaaa

poteps.pep Length: 378 TRANSLATE from: 14 to: 1147

- 1 DEFKDLGLQA CIEHVWRDTI VYLDDDDPIL IGRAYGRVSR HLLHEELLKR
- 51 CVEAGVLYLN SKVDRIVEAT NGHSLVECEG DVVIPCRFVT VASGAASGKF
- 101 LQYELGGPRV SVQTAYGVEV EVDNNPFDPS LMVFMDYRDY VRHDAQSLEA
- 151 KYPTFLYAMP MSPTRVFFEE TCLASKDAMP FDLLKKKLML RLNTLGVRIK
- 201 EIYEEEWSYI PYGGSLPNTE QKTLAFGAAA SMVHPATGYS VVRSLSEAPK
- 251 CAFVLANILR ONHSKNMLTS SSTPSISTQA WNTLWPQERK RORSFFLFGL
- 301 ALILQLDIEG IRSFFRAFFR VPKWMWQGFL GSSLSXADLM LFAFYMFIIA
- 351 PNDMRRGLIR HLLSDPTGAT LIRTYLTF*

FIG. 15B

Chimeric lettuce/potato lycopene $\epsilon\text{-cyclase}$: converts lycopene to $\delta\text{-}$ carotene, the lettuce cDNA converts lycopene to $\epsilon\text{-}$ carotene and the potato cDNA does not produce an active enzyme

(amino acids in lower case are from lettuce and those in uppercase are from the potato cDNA; an $Ava\Pi$ site in common to the two cDNAs was used to construct the chimera)

1	mecfgarnmt	atmavftcpt	ftdcnirhkf	sllkqrrftn	lsassslrqi
51	kcsaksdrcv	vdkqgisvac	eedyvkaggs	elttvamart	Ksmesqskis
101	eklagipign	cildlyvigo	gpaglalaae	sakiginvgi	1gpa/pttnn
151	vavwadefia	laleaciehs	wkdtlvyldd	adpirigray	grvhrdline
201	ellrrcvesa	vsvlsskver	iteapngysl	iecegnitip	criatvasga
251	asokflevel	aGPRVSVQTA	YGVEVEVDNN	PFDPSLMVFM	DYRDYVRHDA
301	OSLEAKYPTE	LYAMPMSPTR	VFFEETCLAS	KDAMPFULLK	KKLMLKLITT
351	GVRTKFTYEE	EWSYIPVGGS	LPNTEQKTLA	FGAAASMVHP	ATGYSVVRSL
401	SEAPKCAEVI	ANTLRONHSK	NMLTSSSTPS	ISTQAWNILW	PQERKKQK2F
451	FLFGLALILO	LDIEGIRSFF	RAFFRVPKWM	WQGFLGSSLS	XADLMLFAFY
501	MFIIAPNDMR	RGLIRHLLSD	PTGATLIRTY	LTF*	

FIG. 16

GAP comparison of Arabidopsis ϵ -cyclase x potato ϵ -cyclase (partial) blosum62.cmp Gap Weight: 12 Average Match: 2.912 Length Weight: 4 Average Mismatch: -2.003 Quality: 1485 Length: 529 Ratio: 3.929 Gaps: 1 Percent Similarity: 79.893 Percent Identity: 76.139 Match display thresholds for the alignment(s): $ = IDENTITY : = 2 . = 1$
151 EDEFNDLGLQKCIEHVWRETIVYLDDDKPITIGRAYGRVSRRLLHEELLR 200
201 RCVESGVSYLSSKVDSITEASDGLRLVACDDNNVIPCRLATVASGAASGK 250
251 LLQYEVGGPRVCVQTAYGVEVEVENSPYDPDQMVFMDYRDYTNEKVRSLE 300
301 AEYPTFLYAMPMTKSRLFFEETCLASKDVMPFDLLKTKLMLRLDTLGIRİ 350 . . 150 AKYPTFLYAMPMSPTRVFFEETCLASKDAMPFDLLKKKLMLRLNTLGVRI 199
351 LKTYEEEWSYIPVGGSLPNTEQKNLAFGAAASMVHPATGYSVVRSLSEAP 400 .
401 KYASVIAEILREETTKQINSNISRQAWDTLWPPERKRQRAFFLFG 445 : : .
446 LALIVOFDTEGIRSFFRTFFRLPKWMWQGFLGSTLTSGDLVLFALYMFVI 495
496 SPNNLRKGLINHLISDPTGATMIKTYLKV 524 . : : : : 350 APNDMRRGLIRHLLSDPTGATLIRTYLTF 378

FIG. 17A

Adonis palaestina Ipil attcatcttc agcagcgctg tcgtactctt tctatatctt cttccatcac taacagtagt cgccgacggt tgaatcggct attcgcctca acgtcaacta 51 tgggtgaagt cactgatgct ggaatggatg ctgttcagaa gcggctcatg 101 ttcgacgacg aatgtatttt ggtggatgag aatgacaagg tcgtcgggca 151 tgattccaaa tacaactgtc atttgatgga aaagatagag gcagaaaatt 201 tgcttcacag agccttcagt gttttcttgt tcaactcaaa atatgaattg 251 cttcttcagc aacgatccgc cacaaaggta acattcccgc tcgtatggac 301 aaacacatgt tgcagtcatc ctctctttcg tgattccgag ctcatagaag 351 aaaattatct cggtgtacga aacgctgcac aaagaaagct tttagacgag 401 ctaggcattc cagctgaaga tgtcccagtt gatgaattta ctcctcttgg 451 tcgcattctt tacaaagctc catctgacgg caaatgggga gagcacgaat 501 tggactatct cctatttatt gtccgagatg tgaaatacga tccaaaccca 551 gatgaagttg ctgatgctaa gtatgttaat cgcgaggagt tgagagagat 601 actgagaaaa gctgatgctg gtgaagaggg actcaagttg tctccttggt 651 ttagattggt tgttgataac tttttgttca agtggtggga tcatgtagag 701 751 cagggtacga ttaaggaagt tgctgacatg aaaactatcc acaagttgac ttaagaggac ttctctcctc tgttctacta tttgtttttt gctacaataa 801 gtgggtggtg ataagcagtt tttctgtttt ctttaattta tggcttttga 851 attigccicg atgtigaact tgtaacatat ttagacaaat atgagacctt 901 qtaagttgaa tttgaggctg aatttatatt tttgggaaca taataatgtt 951 1001 aa

24/45

FIG. 17B

Adonis palaestina Ipi2 1 ttttaaagct ctttcgctcc accaccatca aagccagcca aatttctctg tacaaaagtt aaaaacaccg ctttgggctt tggcccctcc atatcggaat 101 ccttgtttac gatacgcatc taaaccagta attctcggtt ttaatttgtt 151 tcctaaatta ggcccctttc cggaatcccg agaattatgt cgtcgatcag 201 gattaatcct ttatatagta tcttctccac caccactaaa acattatcag 251 cttcgtgttc ttctcccgct gttcatcttc agcagcgttg tcgtactctt tctatttctt cttccatcac taacagtcct cgccgagggt tgaatcggct 301 351 gttcgcctca acgtcgacta tgggtgaagt cgctgatgct ggtatggatg 401 ccgtccagaa gcggcttatg ttcgacgatg aatgtatttt ggtggatgag 451 aatgacaagg tcgtcggaca tgattccaaa tacaactgtc atttgatgga 501 aaagatagag gcagaaaact tgcttcacag agccttcagt gttttcttat 551 tcaactcaaa atacgagttg cttcttcagc aacgatctgc aacgaaggta 601 acattoccyc togtatggac aaacacctgt tgcagccatc coctottccg 651 tgattccgaa ctcatagaag aaaattttct cggggtacga aacgctgcac 701 aaaggaagct tttagacgag ctaggcattc cagctgaaga cgtaccagtt 751 gatgaattca ctcctcttgg tcgcattctt tacaaagctc catctgacgg 801 aaaatgggga gagcacgaac tggactatct tctgtttatt gtccgagatg 851 tgaaatacga tccaaaccca gatgaagttg ctgacgctaa gtacgttaat 901 cqcqaqqaqt tqaaagagat actgagaaaa gctgatgcag gtqaaqaggq 951 aataaagttg tctccttggt ttagattggt tgtggataac tttttgttca 1001 aqtqqtqqqa tcatgtagag gaggggaaga ttaaggacgt cgccgacatg 1051 aaaactatcc acaagttgac ttaagagaaa gtctcttaag ttctactatt tggtttttgc ttcaataagt ggatggtgat gagcagtttt tatgcttcct 1101 ttaattttgg cttttcaatt tgctttatgt gttgaacttg taacatattt 1151 1201 agtcaaatat gagaccttgt gagttgaatt tgaggttata tttatagttt 1251 tgggaacata aaaaaaaaa

25/45

FIG. 18A

Haematococcus pluvialis Ipil 1 ctcggtagct ggccacaatc gctatttgga acctggcccg gcggcagtcc 51 gatgccgcga tgcttcgttc gttgctcaga ggcctcacgc atatcccccg 101 cgtgaactcc gcccagcagc ccagctgtgc acacgcgcga ctccagttta 151 ageteaggag catgeagatg aegeteatge ageceageat eteageeaat 201 ctgtcgcgcg ccgaggaccg cacagaccac atgaggggtg caagcacctg 251 ggcaggcggg cagtcgcagg atgagctgat gctgaaggac gagtgcatct tggtggatgt tgaggacaac atcacaggcc atgccagcaa gctggaqtqt 301 351 cacaagttcc taccacatca gcctgcaggc ctgctgcacc gggccttctc 401 tqtqttcctq tttgacgatc aggggcgact gctgctgcaa cagcgtgcac 451 gctcaaaaat caccttccca agtgtgtgga cgaacacctg ctgcagccac 501 cctttacatg ggcagacccc agatgaggtg gaccaactaa gccaggtggc 551 cgacggaaca gtacctggcg caaaggctgc tgccatccgc aagttggagc 601 acgagetggg gataceageg caccagetge eggeaagege gtttegette 651 ctcacgcgtt tgcactactg tgccgcggac gtgcagccag ctgcgacaca 701 atcagcgctc tggggcgagc acgaaatgga ctacatcttg ttcatccggg 751 ccaacqtcac cttggcgccc aaccctgacg aggtggacga agtcaggtac 801 qtqacqcaaq aggagctgcg gcagatgatg cagccggaca acgggctgca 851 atggtcgccg tggtttcgca tcatcgccgc gcgcttcctt gagcgttggt 901 gggctgacct ggacgcggcc ctaaacactg acaaacacga ggattgggga 951 acggtgcatc acatcaacga agcgtgaaag cagaagctgc aggatgtgaa 1001 gacacqtcat ggggtggaat tgcgtacttg gcagcttcgt atctcctttt 1051 totgagactg aacctgcagt caggtcccac aaggtcaggt aaaatggctc gataaaatgt accgtcactt tttgtcgcgt atactgaact ccaagaggtc 1101 1151 ааааааааа ааааа

26/45

FIG: 18B

Haematococcus pluvialis Ipi2

tggaacctgg cccggcggca gtccgatgcc gcgatgcttc gttcgttgct 51 cagaggeete aegeatatee egegegtgaa eteegeeeag eageeeaget 101 gtgcacacge gegactecag tttaagetea ggageatgea getgettgee 151 gaggaccgca cagaccacat gaggggtgca agcacctggg caggcgggca 201 gtcgcaggat gagctgatgc tgaaggacga gtgcatctta gtggatgctg 251 acgacaacat cacaggccat gccagcaagc tggagtgcca caaattccta 301 ccacateage etgeaggeet getgeacegg geettetetg tgtteetgtt 351 tgacgaccag gggcgactgc tgctgcaaca gcgtgcacgc tcaaaaatca 401 ccttcccaag tgtgtggacg aacacctgct gcagccaccc tctacatggg 451 cagaccccag atgaggtgga ccaactaagc caggtggccg acggcacagt 501 acctggcgca aaagctgctg ccatccgcaa gttggagcac gagctgggga 551 taccagegea ceagetgeeg geaagegegt ttegetteet caegegtttg 601 cactactgtg ccgcggacgt gcagccggct gcgacacaat cagcgctctg 651 gggcgagcac gagatggact acatcttatt catccgggcc aacgtcacct 701 tggcgcccaa ccctgacgag gtggacgaag tcaggtacgt gacgcaagag 751 gagetgegge agatgatgea geeggacaac gggttgeaat ggtegeegtg 801 gtttcgcatc atcgccgcgc gcttccttga gcgttggtgg gctgacctgg 851 acgcggcct aaacactgac aaacacgagg attggggaac ggtgcatcac 901 atcaacgaag cgtgaaggca gaagctgcag gatgtgaaga cacgtcatgg 951 ggtggaattg cgtacttggc agcttcgtat ctcctttttc tgagactgaa 1001 cctgcagagc tagagtcaat ggtgcatcat attcatcgtc tctcttttgt 1051 tttagactaa tctgtagcta gagtcactga tgaatccttt acaactttca 1101 aaaaaaaaa

FIG. 19A

Lactuca sativa Ipil tgccaaaatg ttgaaatttc ccccttttaa aaccattgct accatgatct cticiccata ticticctic tigcigccic ggaaatctic titcccicca atgccgtctc tcgcagccgc tagtgttttc ctccaccctc tttcgtctgc cgctatgggc gattccagca tggatgctgt ccagcgacgt ctcatgttcg 151 201 atgacgaatg cattttggtg gatgagaatg acaaagtggt tggccatgat actaaataca attgtcattt gatggagaag attgaaaagg gaaatatgct 251 301 acacagagca ttcagtgtgt tcttgttcaa ctcgaaatat gaattactcc 351 ttcagcaacg ttctgcaacc aaggtgactt tccctttggt atggacaaac 401 acgtgttgca gccatccact atacagggag agtgagctta ttgacgaaaa 451 cgcccttggg gtgaggaatg ctgcacagag gaagctcctg gatgaactcg 501 gčatcccťgg agčagatgtť ccggttgatg agtťcactcc attgggtcgc 551 attctataca aggccgcatc ggatggaaag tggggagaac atgaacttga 601 ttacctgctg tttatggtac gtgatgttgg tttggatccg aacccagatg 651 aagtgaaaga tgtaaaatat gtgaaccggg aagagctgaa ggaattggta aggaaggcgg atgctggtga agagggtgtg aagctgtccc cgtggttcaa 701 751 attgattgtc gataatttct tgtttcagtg gtgggatcga ctccataagg 801 gaaccctaac cgaagctatt gatatgaaaa caatccacaa actcacataa 851 aaacactaca ctagtaggag agaggattat atgagatatt tgttatatgt 901 gaaattgaaa ttcagatgaa tgcttgtatt tatttctatt tggacaaact 951 tcaacttctt tttgctacct tatcagaaaa aaaaa

FIG. 19B

Lactuca sativa Ipi2

tattcgcttc aaaatctctt ccattaactg ctcaaatctc caccttcgcc ggtcttaatc tccgccggcg cactttcacc accataaccg ccgccatggg 101 151 tgacgattcc ggcatggacg ctgtccagag acgtctcatg tttgatgatg aatgcatttt ggttgafgaa aatgacaatg ttcttgggca tgafaccaaa 201 251 301 tacaattgtc acttgatgga gaagattgag aaagataatt tgcttcatag agcattcagt gtattttat tcaattcaaa atacgaatta ctccttcagc aaaggtcaga aaccaaggtg acatttcctt tggtatggac aaacacctgt 351 tgcagccatc cactatacag agaatcggag ttaattcccg aaaatqccct 401 tggggtcaga aatgctgcac agaggaagct tctagatgaa ctcggtatcc 451 ctgctgaaga tgttccagtt gatgagttca caactttagg tcgcatgttg 501 tacaaggete catetgatgg aaaatggggt gaacatgaag ttgattacet 551 acteticete gigegigaeg tigeegigaa eecaaaeeet galgaggigg cggacattag atacgtgaac caagaagagt taaaagagtt actaaggaag 601 651 gcggatgcgg gtgaggaggg tttgaaattg tccccatggt ttaggctagt 701 ggtggacaac ttcttgttca aatggtggga tcatgtccaa aaggggacac 751 tcaatgaagc aattgacatg aaaaccattc ataagttgat atgaaaaatg 801 gttaaťatít atggiggtgg tttggagcta ataaittgtg tgitcaagtč tcggtccttc ttittitaac gttittttt tttctttat tgggagtgtt 851 901 taligtgtac tigiaacgta ggcccttigg tiacgctita agagtitaat aaagaaccac cgttaattta aaaaaaaaa aaaaaaaa

FIG. 20

Chlamydomonas reinhardtii Ipil

(Note: the isomerase cDNA probably ends at ca. base 1103; the second half of the cDNA is similar to extensin and other hydroxyproline-rich structural proteins)

ggcacgagct cgagtttgtt ttaccatgac atcgggaatt tggaagcttg aactacctca attactcaag taactcgcgg caacacattt cgcgcgccat 101 cgctgttttc tctgctccag ctaccgagca gcattgcttt agatcgcttt 151 gatgicataa actoccacti atatgagato cagtitoato gagoccaago 201 ccagagcgca acctgtctta agccgcggca gggcgtccat gcgcctcgcg 251 301 351 401 caaagccgtg ctctcgttgc gcgtgtcagc tccgccctgt ggccgggagc aggactitca caggeteaaa gegtigeggi gegaatggeg agitegteaa cctgggaagg cacgggcctg agccaggatg acttcatgca gcgggacgag tgcttggtgg tggacgagca ggaccggctg ctaggcaccg ccaacaagta 451 cgactgccac cgcttcgagg cggccaaggg ccagccctgc ggccgcctgc 501 ačcgcgcctt ctccgtgttč ctgttcagčč ccgacggcčg ačtgčtgctg 551 cagcagcgcg cagccagcaa ggtgacgttc ccgggtgtgt ggaccaacac 601 651 701 751 ctgctgctcg cacccgctgg cgggccaggc gccggacgag gtggacctgc cggcggcggt agcctcgggc caggtgccgg gcatcaaggc ggcggcggtg cgcaagctgc agcacgagct ggggataccg ccggagcagg ttcccgcctc ctccttctcc ttcctcacgc gtctgcacta ctgcgccgcc gacaccgcca 801 cgcacggccc ggcggcgag tggggcgagc acgaggtgga ctacgtgctg 851 901 ttcgtgcggc cgcagcagcc cgtcagcctg cagcccaacc cagacgaggt ggacgccacg cgctacgtga cgctgccgga gcttcagtcc atgatggcgg 951 1001 accocggect cagetggage ecctggttee geateetgge caeacageee gccttcctgc ccgcctggtg gggcgacctg aagcggcgct ggcgcccggg cggcagccga ctgtaggact ggggcaccat ccaccgcgtc atgtgaagaa 1051 1101 aaaggggaag caggggcggg agcgggggat gaatgggaat gtgaatgcga 1151 ttgtgatgcg gcgtgggatg aggtctgaag acagggggaa aatcgggggg cgggcgtgag cgtgtgtgta cgtgagcgac aaagccggga ggcggaccgc 1201 1251 1301 gcgatgggta catgtgtgtg cggagggtcg gtgggtcggt cggttgcgcg gcatagcgtg ttgtgtgtgt gcggctgcgc gggtatgtgg gcacccgggc 1351 acggaggaga aggcacacgc aggtggcgcg gaggtgtgtc aggggccatg 1401 ggčgggčctc ačtcctggtc gtgcččagtg gtčtcgtggg cagagtggca 1451 ggggctgcac ccatatgagc ggcgcactgc cgcgctgggc taagtcctta 1501 tcacttggtg aggtgggcg aggtggctgt gggcggcggg cgcagtggca gaaggacacg gtgtgtgagc ggtggagctc tggccgtgcc ggccgtgagg ggcggatagc gatatgacgt tgtgcttggc cgctgtaatg cgggagaatg 1551 1601 1651 1701 tgcaggccgc gagaagcggg cggtggcagg aggccgcagg ctgcagcacc cgttggggag gtgccgcctg caggcgcggc gccgggcggg cctgagtaat 1751 gggcgcctga gtagtggcgg ccacaggagg cgcaggaggc agcagcagga ğğacğagctg ğagğgacccğ ttggcaaccc aaggttgcgc gtgtaacata 1801 1851 gtggccatac aaaaaaaaa aaaa

FIG. 21A

Tagetes erecta Ipil

51 101 151 201 251 301 451 551 561 751 851 851 901	attgtcactt ttcagcgttt gtctgcaacc gccatccact gtaagaaatgt tgaagatgtt aggctccatc ttcatagtga tatcaaatat atgcggggga gataacttct tgaagcaatt tcaaccgaaa caattgttt ttcttgtctt	tggatgctgt gatgagtgtg gatgagagag ttctattcaa aaggtgacat ctacagagaa ctgcacagag cccgttgatc tgatggaaag gagacgttgc gtganccang ggagggtttg tgttcaagtg gatatgaaaa agttcaagtc ttttttaa ttgtgtgtaa	tcagcgacgt acaatgtggt attgaaacag ttcaaaatac ttcctttagt tccgagcttg gaagctgttg agtttactcc tgggggagaac tgtaaacccg aagagttaaa aagctgtctc gtggggatcat ccatacacaa taataattcg gaagttttaa gaagttttaa gaagttttaa gaagttttaa gaagttttaa gaagttttaa gaagttttaa gaagttttaa	ctcatgttfg gggacatgat gtaaaatgct gagttacttc atggaccaac ttcccgaaaa gatgaactcg tttaggtcgc atgaacttga aacccagatg ggagctgcta catggttcag gtgcaaaaagg gctgatatag gctgatatag gctgatatttg ttcgtttcag	acgatgaatg accaaataca gcacagagca ttcagcaacg acctgttgca cgcccttgga gtatccctgc atgctctaca ctacctactt aagtggcgga aggaaagcag gttagtggtt gtacactcac ggtctaccat agcatgttga ttgtaataat
901 951	gaaccattga aaaaaa	tggtttgcaa	tttcaagttc	ctatcgacat	gtagtgatct

FIG. 21B

Oryza -	sative Ipil				
1	cctccctttg	cctcgcgcag	aggcggccgc		ccgcgaggat
51	ggccggcgcc	gccgccgccg	tggaggacgc	cgggatggac	gaggtccaga
101	agcggctcat	gttcgacgac		tggtggatga	acaagacaat
151	gttgttggcc	atgaatcaaa		catctgatgg	aaaaaatcga
201		ctacttcata		tgtattcctg	ttcaactcaa
251		cctactccag		caacaaaggt	tacatttcct
301	ctagtttgga	ccaacacttg		cctctgtacc	gtgagtctga
351		gaaaactacc		aaatgctgct	cagaggaagc
401	tcttggatga			atgtgccagt	tgaccaattc
451		gtcggatgct		ccatctgatg	gaaaatgggg
501	tgaacacgag			cgtccgcgac	gtgaaggtag
551	tcccgaaccc	ggacgaagtg		aatacgtgag	ccgtgagcag
601	ctgaaggagc	tcatccgcaa	agcggacgcc	ggagaggaag	gcctgaagct
651	gtctccctgg	ttccggctgg	ttgttgacaa	cttcctcatg	ggctggtggg
701	atcacgtcga		ctcaacgagg	ccgtggacat	ggagaccatc
751	cacaagctga	agtaaggact	gcgatgttgt	ggctggaaag	aatgatcctg
801	aagactctgt	tcttgtgctg	ctgcatatta	ctcttaccag	ggaagttgca
851	gaagtcagaa	gaagettttg	tatgtttctg	ggtttggagc	ttggaagtgt
901 951	tgggctctgc	tgactgagag	attcccttat	agagtgtcta	tgttaattta
1001	gcaaacttct agaacaatag	atattataca catgttccat	tgattagttagttagttatttgc	attgttcggt	gtctgaataa
TOOT	ayaacaatay	catguittat	geetattige	L	

The transmission of the tr

The first state of the first that the

		_				- 10	_	0,	_	10	_	0	6	~	0	6	2	-	♥
27	75	7 26	55	2 8	29	88	8		107	155	107	170	109	Ξ	16	Ò	16	1	174
16 60 61 75 76 90 90	MLKFPPF KTIATMISSPYSSFL LPRKSSFPPMPSLAAASVFLJHPLSSA AMGDSS-MDAVQR RLMFDDECILVDEND	NSPRRGLNRLFASTS TWGEVADAGMDAVQK RLMFDDECTLVDEND		PRKLPNFRAFSGT AMTDTKDAGMDAVQR RLMFEDECILVDETD		MLRSLLRGLTHI PRVNSAQQPSCAHAR LQFKLRSMQLLSEDRIDH MRGASIWAG-GQSQD ELMLKDECILVOVED MIDSELIDGITUT DDVNSAQDSCAHAR LOFKIRSMANIMOP SISANISRAFDRIDH MRGASIWAG-GOSOD ELMLKDECILVOVED		136 150 151 165 166 180	KIETGKMLHRAFS VFLENSKYELLLQQR SATKVTFPLVWTNTC CSHPLYRESELVPENALGVRNA	(IE KGHMLHRAFS VFLENSKYELLLQQR SATKVIFPLWATNIC CSHPLYRES ELIDENALGVRNA	(IE KONI LHRAFS VFLFNSKYELLLQQR SETKVTFPLVWTNTC CSHPLYRES ELIPENALGVRNA	KIEAEHLLHRAFS VFLFNSKYELLLQQR SATKVTFPLVWINIC CSHPLFROSELIEENFLGVRNA	CSHPLFRDS	1 1 1	1				AAKGQPCGRLHRAFS VELFSPDGRFLLQQR AASKVTFPGVWTNTC CSHPLAGQAPDEVDL PAAVASGQVPGTKAA
45 46	Sdl	118881				LLS	RVSSAL N	135 136	LLLOOR	LLLOOR	LLLOOR	LLLQQR	LLLQQR	LLLOOR	LLLQQR	LLLQQR	LLLQQR	LLLQQR	11100R
30 31	L LPRKSSFPPM	H I MRCRTI SI	יייייייייייייייייייייייייייייייייייייי	R FAHRPLSS	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	IR LQFKLRSMQL ID 1 DFK1 RSMOW	IR LAQSRALVAR	120 121	S VFLENSKYEI	S VFLENSKYEL	S VFLFNSKYEI	'S VFLFNSKYEI	'S VFLFNSKYEI	S VFLFNSKYEI	S VFLFNSKYEI	S VFLFNSKYE	S VFLFDDQGRI	S VFLFDDQGRI	FS VFL FSPDGRI
	MLKFPPF KTIAIMISSPYSSFI	TISSELE TIME TO THE TOTAL TO TH	MOSIKIMPLISIFSI IINILSASCASIANI	KVCSCI FNI PI I RI RSI AI SSSESSFR FAHRPLSSIS		MLRSLLRGLTHI PRVNSAQQPSCAHAR LQFKLRSMQLLS	mrssfiepk praqpylsrgrasm	105 106				_		NVVGHESKYNCHLME KIESENLLHRAF	RVVGHVSKYNCHLME NIEAKHLLHRAF	RVVGHDTKYNCHLME KIEAENLLHRAF			RLLGTANKYDCHRFE AAKGQPCGRLHRAF
These amino acid sequences were present	1 Terecta 1	3 L.sativa 2	4 A.palaestina 2 5 A.palaestina 1	6 O.sativa 1	8 A. thal iang 2	9 H.pluvialis 1	10 H.pluVlatis 2 11 C.reinhardtii 1		1 Torocta 1	2 1 sativa 1	3 I sativa 2	4 A.palaestina 2	5 A.palaestina 1	6 O.sativa 1	7 A.thaliana 1	8 A. thaliona 2	9 H.pluvialis 1	10 H.pluvialis 2	11 C.reinhardtii 1

F16. 22 A

188 236 188 251 190 194 241 247 259	
226 240 241 255 256 270 EHELDYLLFIVRD VAVNPNPDEVADIKY VSHEELKELLRKADA EHELDYLLFIVRD VGLDPNPDEVADIKY VNREELKELVRKADA EHELDYLLFIVRD VAVNPNPDEVADIRY VNREELKELLRKADA EHELDYLLFIVRD VKYDPNPDEVADAKY VNREELKEILRKADA EHELDYLLFIVRD VKYDPNPDEVADAKY VNREELKEILRKADA EHELDYLLFIVRD VKYOPNPDEVADIKY VSREELKELVKKADA EHELDYLLFIVRD VKLQPNPDEVAEIKY VSREELKELVKKADA EHEWDYLLFIVRD VKLQPNPDEVAEIKY VSREELKELVKKADA EHEWDYLLFIRAN VILAPNPDEVDEVRY VIQEELRQMAQP EHEMDYILFIRAN VILAPNPDEVDEVRY VIQEELRQMAQP EHEWDYLLFIRAN VILAPNPDEVDEVRY VIQEELRQMAQP	Tagetes erecta (marigold) Lactuca sativa (romaine lettuce) Lactuca sativa (romaine lettuce) Adonis palaestina (pheasant's eye) Oryza sativa (rice) Arabidopsis thaliana Haematococcus pluvialis Chlamydomonas reinhardtii
240 241 IVRD VAV WWRD VGL LVRD VKY IVRD VKY IVRD VKY IVRD VKY IVRD VKV IVRD VKV IVRD VKV IVRD VKV	232 280 229 234 238 284 233 305
AQRKLLDELGIPAED VPVDQFTPLGRMLYKAPSDGKWG AQRKLLDELGIPAED VPVDEFTPLGRILYKAPSDGKWG AQRKLLDELGIPAED VPVDEFTPLGRILYKAPSDGKWG AQRKLLDELGIPAED VPVDEFTPLGRILYKAPSDGKWG AQRKLLDELGIPAED VPVDEFTPLGRILYKAPSDGKWG AQRKLLDELGIPAED VPVDEFTPLGRMLYKAPSDGKWG AQRKLLDELGIVAED VPVDEFTPLGRMLYKAPSDGKWG AQRKLEHELGIPAED VPVDEFTPLGRMLYKAPSDGKWG AIRKLEHELGIPAHQ LPASAFRFLTRLHYC AADVQPAATQSALWG ANRKLEHELGIPAHQ LPASAFRFLTRLHYC AADVQPAATQSALWG AVRKLQHELGIPPEQ VPASSFSFLTRLHYC AADVQPAATQSALWG	GEEGLKLSPWFRLVV DNFLFKAMDHVQK GTLTEAIDMKTI HKLIGEEGVKLSPWFRLVV DNFLFGAMDRLHK GTLTEAIDMKTI HKLIGEEGVKLSPWFRLIV DNFLFKAMDHVQK GTLNEAIDMKTI HKTTGEEGLKLSPWFRLVV DNFLFKAMDHVEG GTLNEAIDMKTI HLTGEEGLKLSPWFRLVV DNFLFKAMDHVEG GTIKEVADMKTI HKLTGEEGLKLSPWFRLVV DNFLMKAMDHVEK GTLNEAVDMFTI HKLKGEEGLKLSPWFRLVV DNFLMKAMDHVEK GTLNEAVDMFTI HKLKDHGLQWSPWFRIJA ARFLERWADLDA ALNTDKHEDMGTV HHINEADNGLQWSPWFRIJA ARFLERWADLDA ALNTDKHEDMGTV HHINEADPGLSWSPWFRIJA TQPAFLPAWMGDLKR RWRPGGSRLSDWGTV HHINEADPGLSWSPWFRILA TQPAFLPAWMGDLKR RWRPGGSRLSDWGTV HRINEADPGLSWSPWFRILA TQPAFLPAWWGDLKR RWRPGGSRLSDWGTV HRINEADPGLSWSPWFRILA TQPAFLPAWWGDLKR RWRPGGSRLSDWGTV HRINEA
1 T. erecta 1 2 L. sativa 1 3 L. sativa 2 4 A. palaestina 2 5 A. palaestina 1 6 O. sativa 1 7 A. thaliana 1 8 A. thaliana 2 9 H. pluvialis 1 10 H. pluvialis 2	1 T.erecta 1 2 L.sativa 1 3 L.sativa 2 4 A.palaestina 2 5 A.palaestina 1 6 O.sativa 1 7 A.thaliana 1 8 A.thaliana 2 9 H.pluvialis 1 10 H.pluvialis 2 11 C.reinhardtii 1

F16.22B




FIG. 23A

09/701395

```
Comparison using GAP program of the Genetics Computer Group
                             Average match:
                                          10.000
                    50
       Gap Weight:
                                           0.000
     Length Weight:
                           Average Mismatch:
                 17392
                                   Length:
                                            1904
         Quality:
           Ratio: 9.411
                                     Gaps:
                          Percent Identity:
 Percent Similarity: 95.331
Match display thresholds for the alignment(s):
                       : = 5
             = IDENTITY
Adonis palaestina E-cyclase #3 x Adonis palaestina E-cyclase #5
   1 gagagaaaagagtgttatattaatgttactgtcgcattcttgcaacac. 49
         .aaaggagtgttctattaatgttactgtcgcattcttgcaacact 44
  50 .atattcagactccattttcttgttttctcttcaaaacaacaactaatg 98
  99 tga.cggagtatctagctatggaactacttggtgttcgcaacctcatctc 147
  95 tgagcagagtatctggctatggaactacttggtgttcgcaacctcatctc 144
 148 ttcttgcctgtctggacttttggaacaagaaaccttagtagttcaaaac 197
     145 ttcttgccctgtgtggacttttggaacaagaaaccttagtagttcaaaac 194
 198 tagcttataacatacatcgatatggttcttcttgtagagtagattttcaa 247
     248 gtgagggctgatggtggaagcgggagtagaacttctgttgcttataaaga 297
     245 gtgagagctgatggtggaagcgggagtagaagttctgttgcttataaaga 294
  298 gggttttgtggacgaggaggattttatcaaagctggtggttctgagcttt 347
  295 gggttttgtggatgaagaggattttatcaaagctggtggttctgagcttt 344
  348 tgtttgtccaaatgcagcaaacaaagtctatggagaaacaggccaagctc 397
  345 tgtttgtccaaatgcagcaaacaaagtctatggagaaacaggccaagctc 394
```

FIG. 23B

398 gccgataagitgccaccaaiacctttcggagaatctgtgatggacttggt	447
395 gccgataagttgccaccaataccttttggagaatccgtgatggacttggt	444
448 tgtaataggttgtggacctgctggtctttcactggctgcagaagctgcta	497
445 tgtaataggttgtggacctgctggtctttcactggctgcagaagctgcta	494
498 agctaggcttgaaagttggccttattggtcctgatcttccttttacaaat	547
495 agctagggttgaaagttggccttattggtcctgatcttccttttacaaat	544
548 aattatggtgtgtgggaagacgagttcaaagatcttggacttgaacgttg	597
545 aattatggtgtgtgggaagacgagttcaaagatcttggacttgaacgttg	594
598 tatcgagcatgcttggaaggacaccatcgtatatcttgacaatgatgctc	647
595 tatcgagcatgcttggaaggacaccatcgtatatcttgataatgatgctc	
648 ctgtccttattggtcgtgcatatggacgagttagccggcatttgctgcat	697 694
645 ctgtccttattggtcgtgcatatggacgagttagtcgacatttgctacat 698 gaagagttgctgaaaaggtgtgtcgagtcaggtgtatcatatctgaattc	
695 gaggagttgctgaaaaggtgtgtggagtcaggtgtatcatatctggattc	744
748 taaagtggaaaggatcactgaagctggtgatggccatagtcttgtagttt	797
745 taaagtggaaaggatcactgaagctggtgatggccatagccttgtagttt	794
798 gtgaaaacgacatctttatcccttgcaggcttgctactgttgcatctgga	847
795 gtgadatgagatetttateeettgeaggettgetaetgttgeatetgga	
848 gcagcttcagggaaacttttggagtatgaagtaggtggccctcgtgtttg	
845 gcagcttcagggaaacttttggagtatgaagtaggtggccctcgtgtttg	
898 tgtccaaactgcttatggtgtggaggttgaggtggagaacaatccatacg	
895 tgtccaaaccgcttatggggtggaggttgaggtggagaacaatccatacg	J 1 T

FIG. 23C

948	atcccaacttaatggtatttatggactacagagactatatgcaacagaaa	997
945	atcccaacttaatggtattcatggactacagagactatatgcaacagaaa	994
998	ttacagtgctcggaagaagaatatccaacatttctctatgtcatgcccat	1047
995	ttacagtgctcggaagaagaatatccaacatttctctatgtcatgcccat	1044
1048	gtcgccaacaagacttttttttgaggaaacctgtttggcctcaaaagatg	1097
1045	gtcgccaacaagacttttttttgaggaaacctgtttggcctcaaaagatg	1094
1098	ccatgcctttcgatctactgaagagaaaactaatgtcacgattgaagact	1147
1095	ccatgccattcgatctactgaagagaaaactgatgtcacgattgaagact	1144
1148	ctgggtatccaagttacaaaaatttatgaagaggaatggtcttatattcc	1197
	ctgggtatccaagttacaaaagtttatgaagaggaatggtcatatattcc	1194
	tgttgggggttctttaccaaacacagagcaaaagaacctagcatttggtg	1247
	tgttggtggttctttaccaaacacagagcaaaagaacctagcatttggtg	1244
	ctgcagcaagcatggtgcatccagcaacaggctattcggttgtacgatca	1297 1294
	ctácaácaáácatáátgcatccagcaacagactattcggttgtacggtca ctatcagaaagctccaaaatatgcttctgtaattgcaaagattttgaagca	1347
		1344
	agataactcigcatatgtggtttctggacaaagcagtgcagt	1397
		1394
1398	· · · · · · · · · · · · · · · · · · ·	1447
1395		1444
1448	gcattctttcttttcgggttagagcttattgtgcagctagatattgaagc	1497
144!	5 gcattctttctttttggattagagcttattgtgcagctagatattgaagc	1494

1498	aaccagaacgttctttagaaccttcttccgcttgccaacttggatgtggt	1547
1495	aaccagaacattctttagaaccttcttccgcttgccaacttggatgtggt	1544
1548	ggggtttccttgggtcttcactatcatctttcgatcttgtattgttttcc	1597
1545	ggggtttccttgggtcttcactatcatctttcgatctcgtcttgttttcc	1594
1598	atgtacatgtttgttttggccccgaacagcatgaggatgtcacttgtgag	1647
1595	atgtacatgtttgttttggcgccaaacagcatgaggatgtcacttgtgag	1644
1648	acatttgctttcagatccttctggtgcagttatggttaaagcttacctcg	1697
1645	acatttgctttcagatccttctggtgcagttatggtaagagcttacctcg	1694
1698	aaaggtaatctgttttatgaaactatagtgtctcattaaataaatga	1744
	àààggtàgtctcatctattàttàáàctctàgtgtttcàccàààtàààtgà	1744
	ggatccttcgtatatgtatatgatcatctctatgtatatcctatattcta	1794
	ggátccttcgaatgtgtatatgatcatctctatgtatatcctgtactcta	1794
	atctcataaagtaatcgaaaattcattgatagaaaaaaaa	1844
	atctcataaagtaaatgccgggtttgatattgttgtgtcaaaccggccaa	1844
	aaaa	1848
1845	tgatataaagtaaatttattgatacaaaagtagtttttttt	1894

FIG. 23D

GAP program of Genetics Computer Group blosum62.cmp
Gap Weight: 12 Average Match: 2.912 Length Weight: 4 Average Mismatch: -2.003 Quality: 2728 Length: 530 Ratio: 5,147 Gaps: 0 Percent Similarity: 99,623 Percent Identity: 99.057 Match display thresholds for the alignment(s): = IDENTITY := 2 . = 1
Adonis palaestina ε-cyclase #3 x Adonis palaestina ε-cyclase #5
1 MELLGVRNLİSSCPVWTFGTRNLSSSKLAYNIHRYGSSCRVDFQVRADGG 50
51 SGSRTSVAYKEGFVDEEDFIKAGGSELLFVQMQQTKSMEKQAKLADKLPP 100
101 IPFGESVMDLVVIGCGPAGLSLAAEAAKLGLKVGLIGPDLPFTNNYGVWE 150
151 DEFKDLGLERCIEHAWKDTIVYLDNDAPVLIGRAYGRVSRHLLHEELLKR 200
201 CVESGVSYLNSKVERITEAGDGHSLVVCENDIFIPCRLATVASGAASGKL 250
251 LEYEVGGPRVCVQTAYGVEVEVENNPYDPNLMVFMDYRDYMQQKLQCSEE 300
301 EYPTFLYVMPMSPTRLFFEETCLASKDAMPFDLLKRKLMSRLKTLGIQVT 350

38/45

351	KIYEEEWSYIPVGGSLPNTEQKNLAFGAAASMVHPATGYSVVRSLSEAPK	400
351	:	400
401	YASVIAKILKQDNSAYVVSGQSSAVNISMQAWSSLWPKERKRQRAFFLFG	450
401		450
451	LELIVOLDIEATRTFFRTFRLPTWMWWGFLGSSLSSFDLVLFSMYMFVL	500
451	LELIVQLDIEATRTFFRTFFRLPTWMWWGFLGSSLSSFDLVLFSMYMFVL	500
501	APNSMRMSLVRHLLSDPSGAVMVKAYLER* 530	
501		

FIG. 24B

WO	99/63055	.39/	,45	r C1/0599/12121
1			•	09/701395
			454400=	
•	103 102 102 100 100 100 75 75 75 75 75 75 75 75 75 75 75 75 75	38.	205 204 204 209 209 202	181 184 184 187 188 188 189
* }		 	FE ES ES ES ES ES ES ES ES ES ES ES ES ES	SACACAN
;	TOPE THE PERSON OF THE PERSON	- E	\$15 5 5 5 5 5 5 5 5 7 1 1 1 1 1 1 1 1 1 1	<u> </u>
	K K K K K K K K K K K K K K K K K K K	X XX		
100	QAKI QAKI QAKI QSKI QSSI PETI PETI PETI PETI	PEI	* <u> </u>	X
	MDE MEK MES MES MDQ MDQ MDQ MDQ MDQ MDQ MDQ MES MDA MDQ MDQ MES MDQ MDQ MDQ MDQ MDQ MDQ MDQ MDQ MDQ MDQ	ELV	<u> </u>	**************************************
	ONKE OTKS OKKE ONKS SALI SALI	SALI		A A C C C C C C
*	GMQ GMQ GMQ GMQ GMQ GMQ GMQ GMQ GMQ GMQ	ass ass svss	GGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	28 P V V V V V V V V V V V V V V V V V V
,	LLEV LLEV LVEV LLEV SSSVV SSSVV VCVK VCVK	YCV GYI(LIGORIAN TERMINATION TO THE PROPERTY OF THE PR	KOLL KOLL KOLL KOLL KOLL KOLL
	GRS CRN	X X	DODP DOKP TOAP TOAP DOEP	EGVK ONSK ONTA ONTA ONTT
8	VKAG VKAG IKAG VKAG VKAG VKAG	YOK		
	EDF EDF EDY EDY KIKI SRNI SRNI SRNI EDY	SK KS	7110 7110 7110 7110 7110	Savv Savv Stvv Savv Galv
		RFG! KFF.	NAME OF THE PARTY	S S S S S S S S S S S S S S S S S S S
*	105- 105- 105- 105- 105- 105- 100- 100-	9 6 A	AGIEHV AGIEHV AGIEHA AGIEHA AGIEHA AGIEHA	
	AVRE VOK VOK VVK SRVF KLQF KRQF KRQF	S S S S S S S S S S S S S S S S S S S		
0	ESCV RSSV RTSV CV DSCV DSCV NPYH NPYH VSTS FSSV FFRS	SPSS TSPS	XOLC XOLC XOLC XOLC XOLC XOLC XOLC	
9	SYRNIRFGL-CSVRASGGGSSGSESCVAVREDFADEEDFVKAGGSE1LFVQMQQIKSMEKQAKLADKÜPPIS AYNIHRYGSSCRVDFQVRADGGSGSRSSVAYKEGFVDEEDFIKAGGSELLFVQMQQIKSMEKQAKLADKÜPPIP AYNIHRYGSSCRVDFQVRADGGSGSRTSVAYKEGFVDEEDFIKAGGSELLFVQMQQIKSMEKQAKLADKÜPPIP QRRFTNLSA-SSSLRQIKCSAKSDRCVVDKQGISVADEEDFYKAGGSELFVQMQQIKSMEKQAKLSDEÜPPIP QRRFTNLSA-SSSLRQIKCSAKSDRCVVDKQGISVADEEDFYKAGGSELFVQMQRIKSMESQSKLSEKÄÄQIP QEKSIFLAY-EQYESKCNSSSGSDSCVVDKEDFADEEDFYKAGGSELLFVQMQRKXMDAQSSLSQKÄPPIP LKTPNKLDFFIPQFHGFERLCSNNPYHSRVRLGVKKRAIKIVSSVVSGSAALLDLVPETKKENÄDFEL LRTHNKLELLPTLHGFAEKQHLVSTSKLQNQVFRIASRNIHPCRNGTVKAGSSALLELVPETKKENÄDFEL LKTPNNLEFLHGFGAVVSAFSSVKSQKFGAKKFCEGLGSRSVCVKASSSALLELVPETKKENÄDFEL LKTPNNLEFLNPHHGFAVVSAFSSVKSQKFGAKKFCETLGRSVCVKGSSSALLELVPETKKENÄDFEL LKTPNNLEFLNPHHGFAVVSAFSSVKSQKFGAKKFCETLGRSVCVKASSSALLELVPETKKENÄDFEL	LRTYNSFEFVHPSNKFAGNLNNUNQLNQSKSQFQDFRFGPKKSQFKLQQKYCVKASSSALLELVPETKKENLDFDL LRTYNRLELLYPLHELAKRHFLSPSPNPQNPNFKFFSRKPYQKKCRNGYTGVSSNQLLDLVPETKKEHLEFDL	* 160 * 200	LSVCSTOPS - PKL IWPNINYGWWOE FEAMOLL DCLOTTWSGAVYYTÖDNSKKYLDRPYGRVNRKQLKSKMLQKÇITN LSVCSTOPS - PKL IWPNINYGWWOE FEAMOLL DCLOTTWSGAVYYTÖDNSKKYLDRPYGRVNRKQLKSKMLQKÇITN LSVCSTOPN - PKL IWPNINYGWWOE FEAMOLL DCLOATWSGAAVYTÖDNTAKOLHRPYGRVNRKQLKSKMMQKÇITUN LSVCSTOPN - PKL IWPNINYGWWOE FEAMOLL DCLOATWSGAAVYTÖDNTTKOLHRPYGRVNRKQLKSKMMQKÇITUN LSVCSTOPN - PKL IWPNINYGWWOE FEAMOLL DCLOATWSGAVYTÖDNTTKOLORPYGRVNRKQLKSKMMQKÇITUN LTWCSTOPS - PKL IWPNINYGWWOE FEAMOLL DCLOATWSGAVYTÖDNTTKOLORPYARVINRKQLKSKMMQKÇITAN LTWCSTOPN - PKL IWPNINYGWWOE FEAMOLL DCLOATWSGAVYTÖDRSTKNLSRPYARVINRKQLKTKMLQKÇIAN LSVVSTOPN - PKL IWPNINYGWWOE FEAMOLL DCLOATWSGAVYTÖDRSTKNLSRPYARVINRKQLKTKMLQKÇIAN
	ADGGS ADGG CSAK CCNSS CNAA CCNAA CCNAA CCNAA	N N	COVER GOVER	
*	-RAS FQVR FQVR RQ1K RQ1K KQ1K GFG- GFG- GFG-	KFAC KFAC ELA-	PINN PINN PINN PINN PINN PINN PINN PINN	NNA NNA NNA NNA NNA NNA NNA NNA NNA NNA
	SSV- SRVD SSSL SSSL SSSL SY- TLH TPQF	4PSN 4PLH 7PLH		
	GC-C GSS(GSS(GSS(A-SSA-SSA-SSA-SSA-SSA-SSA-SSA-SSA-SSA		* KYGL IGPOLP- KYGL IGPOLP- KYGL IGPOLP- NVGL YGPOLP- NVGL YGPOLP- NVGL YGPOLP-	2
\$	NIRF IHRY IHRY IHRY FTNL SIFL PNNL PNNL	YNS!	* 1000000	
	SYRU AYN QRRU QRRU QEK LKT LKT			
	GTRNLSSSKLA GTRNLSSSKLA GTRNLSSSKLA GTRNLSSSKLA GTRNLSSSKLA GTRNLSSSKLA GTRNLSSSSKLA 	MOTF	AKLG AKLG AKLG AKLG AKLG	SEAG SEAG SEAG SEAG SEAG 3-GG
*	KFPV RNLS RNLS RHKF		140 AESAI AEAAI AESAI AESAI AESAI	
	CRR FGT CNI		LS LS LS LS LS LS LS LS LS LS LS LS LS L	LAVA LAVA LAVA SCS
50	SWS- SWT- WWT- PRFTI		120 * 140 16	PLYDTSKSQVVDLATVĢGGPAGLAVAQQVSEAGI PAYDPSKGVVVDLAVVGGPAGLAVAQQVSEAGI PMYDPSKGVVVDLAVVGGPAGLAVAQQVSEAGI PMYDPSKGLVVDLAVVGGPAGLAVAQQVSEAGI PMYDPSKGLVVDLAVVGGPPAGLAVAQQVSEAGI PMYDPSKNVVVDLVVVGGPSGLAVAQQVSEAGI PLYDPSKALTLDLAVVGGPPLARSCSTSLG-GG
2	SSCF SSCF SSCF FTCF FTCF			\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \
	WMAV IMAV IMAA			
*	#		120 DGAL ESVM ESVM ESVM OTVL	SQVV GLVV GLVV GLVV GLVV GLVV GLVV
	GARN GVRN GVRN GARN GARN			758 758 768 768 768 768 768 768
Ø	MECUGARNE-AAMAVSTFPSWS-CRRKFPVVKRYSYRNIRFGL-CSVRASGGGSSGSESCVAVREDFADEEDFVKAGGSE1LFVQMQQNKDMDEQSKLVDKÜPPIS MELLGVRNLISSCPW4T-FGTRNLSSSKLAYNIHRYGSSCRVDFQVRADGGSGSRTSVAYKEGFVDEEDFIKAGGSELLFVQMQQTKSMEKQAKLADKÜPPIP MELLGVRNLISSCPW4T-FGTRNLSSSKLAYNIHRYGSSCRVDFQVRADGGSGSRTSVAYKEGFVDEEDFIKAGGSELLFVQMQQTKSMEKQAKLADKÜPPIP MECFGARNATATMAVFTCPRFTDCNIRHKFSLLKQRRFTNLSA-SSSLRQIKCSAKSDRCVVDKQGISVADEEDFVKAGGSELLFVQMQQTKSMESQSKLSEKÄÄQIP MECKGARNATATMAVFTCPRFTDCNIRHKFSLLKQRRFTNLSA-SSSLRQIKCSAKSDRCVVDKQGISVADEEDFVKAGGSELLFVQMQQTKDMDQQSKLSEKÄÄQIP MECVGVQNV-GAMAVLTRPRLNRASGGELCQEKSIFLAY-EQYESKCNSSSGSDSCVVDKEDFEEEEDFVKAGGSELLFVQMQQTKDMDQQSKLSDÄPRPP MESMRAG-HATATMAAFTCPRFM		16 76 16 16 86	PAYE PAYE PMYE PMYE PLYE
F16.25A				
נע	PotatoE ArabidopsisE AdonisE1 AdonisE2 LettuceEE TomatoE ArabidopsisB AdonisB PepperB TomatoB	∞	PotatoE ArabidopsisE AdonisE1 AdonisE2 LettuceEE TomatoE	ArabidopsisB AdonisB PepperB TomatoB TobaccoB MarigoldB DaffodilB
9	Arabidops AdonisE1 AdonisE2 LettuceEE TomatoE MarigoldE Arabidops AdonisB PepperB	TobaccoB MarigoldB DaffodilB	PotatoE Arabidops AdonisE1 AdonisE2 LettuceEE TomatoE	Arabidops Arabidops AdonisB PepperB TomatoB TobaccoB MarigoldB DaffodilB
正	Pote Arat Ador Ador Tome Mari Ador Pepi	Tob Mar Daf	Pot Ara Ado Ado Let Iom	Ara Ado Ado Pep Tom Tob Mar Daf

F16. 25


SSTRIFLEETSLVARPGLKMEDIQERMAYRLKHLGIKVKSIEEDERCVIFMGGPLPVLPQRVLGEGDTACMVHPSTGYMVARTIJAAARIVAKSIIRYLNNEKSM---VAD

MarigoldB DaffodilB

Fobacco8

SSNK JFLEETSLYARPGLRYDDIQERWYARLNHLGIKVKSIGEDEHCVIPYGGSLPVIPQRVVGTGBTAGLVHPSJGYMVARTLAAAPVVANAIIHYLGSEKDL----

SSNRI£LEETSJVARPGLKMEDIQERMARLINHLGIRIKSIEEDERCVIPMGGPLPVIPQRVVGIGGTACKVHPSIGYMVARITLAAAPIVANSIVQYLVSDSGL-----



43/45




FIG. 28A

GAP of: Arabidopsis epsilon cyclase to Lettuce epsilon cyclase Gap Weight: Average Match: 2.912 12 4 Average Mismatch: -2.003Length Weight: 534 Length: Quality: 1837 Ratio: 3.499 Gaps: Percent Similarity: 76.381 Percent Identity: 69.905 Match display thresholds for the alignment(s): = IDENTITY : = 2 Arabidopsis x Lettuce 1 MECVGARNF.AAMAVSTFPSW...SCRRKFPVVKRYSYRNIRFGLCSVRA 46 1 MÉCFGARNMTATMÁVFTCPRFTDCNIRHKFSLLKQRRFTNLSASSSLRQI 50 1111:111111: 11111. 1 1: 1111 51 KCSAKSDRCVVDKQGISVÄDEEDYVKÄGGSELFFVQMQRTKSMESQSKLS 100 151 YGVWODEFIGLGLEGCIEHSWKDTLVYLDDADPIRIGRAYGRVHRDLLHE 200 197 ELLRRCVESGVSYLSSKVDSITEASDGLRLVACDDNNAIPCRLATVASGA 246 247 ASGKLLQYEVGGPRVCVQTAYGVEVEVENSPYDPDQMVFMDYRDYTNEKV 296 297 RSLEAEYPTFLYAMPMTKSRLFFEETCLASKDVMPFDLLKTKLMLRLDTL 346

45/45

FIG. 28B

	• • • • • • • • • • • • • • • • • • • •	
347	GIRILKTYEEEWSYIPVGGSLPNTEQKNLAFGAAASMVHPATGYSVVRSL	396
351	:	400
397	SEAPKYASVIAEILREETTKQINSNISRQAWDTLWPPERKRQRAF	441
401		450
442	FLFGLALIVOFDTEGIRSFFRTFFRLPKWMWOGFLGSTLTSGDLVLFALY	491
451		500
492	MFVISPNNLRKGLINHLISDPTGATMIKTYLKV* 525	
501	. : : : : MFVIAPHSLRMELVRHLLSDPTGATMVKAYLTI* 534	

Docket No. 108172-00022

ARENT FOX KINTNER PLOTKIN & KAHN, PLLC

Nikaido, Marmelstein, Murray & Oram Intellectual Property Group

Declaration For U.S. Patent Application

As a below named inventor, I hereby declare that: My residence, post office address and citizenship are as stated below my name. I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled (Insert Title) GENES OF CAROTENOID BIOSYNTHESIS AND METABOLISM AND METHODS OF USE THEREOF the specification of which is attached hereto unless the following box is checked: June 2, 1999 was filed on As PCT International Application PCT/US99/12121 Number and was amended on December 4, 2000 and/or was filed on As U.S. Patent Application and was amended on I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claim(s), as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate or PCT International Application having a filing date before that of the application(s) for which priority is claimed: Priority Claimed ☐ Yes ☐ No (List prior (Number) (Country) (Day/Month/Year Filed) foreign Yes Yes ☐ No applications) (Number) (Country) (Day/Month/Year Filed) ☐ Yes ☐ No (Number) (Country) (Day/Month/Year Filed) I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below. M (Application Number) (Filing Date) (Application Number) (Filing Date) See attached list for additional prior foreign or provisional applications. PL I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s) or §365(c) of any PCT International application(s) designating the United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior application(s) (U.S. or PCT) in the prior application(s) (U.S. or PCT) in the prior application (s) disclosed in the prior application(s) (U.S. or PCT) in the manner provided by the first paragraph of 35, U.S.C. §112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 C.F.R. §1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

And I hereby appoint the firm of Arent Fox, Customer Number 004372 including as principal attorneys: Robert B. Murray, Reg. No. 22,980; Charles M. Marmelstein, Reg. No. 25,895; George E. Oram, Jr., Reg. No. 27,931; Douglas H. Goldhush, Reg. No. 33,125; David T. Nikaido, Reg. No. 22,663; Richard J. Berman, Reg. No. 39,107; King L. Wong, Reg. No. 37,500; James A. Poulos, III, Reg. No. 31,714; Murat Ozgu, Reg. No. 44,275; Robert K. Carpenter, Reg. No. 34,794; Gregory B. Kang, Reg. No. 45,273; Rustan Hill, Reg. No. 37,351; Carl Schaukowitch, Reg. No. 29,211; Kevin Turner, Reg. No. 43,437; Rhonda C. Barton, Reg. No. P47,271 and Hans J. Crosby, Reg. No. 44,634.

June 2, 1998

(Filing Date)

June 2, 1998

(Filing Date)

Please direct all communications to the following address:

09/088,724

09/088,725

(Application Serial No.)

(Application Serial No.)

(List prior U.S.

Applications or

applications

PCT International

designating the US)

ARENT FOX KINTNER PLOTKIN & KAHN, PLLC 1050 Connecticut Avenue, N.W., Suite 600 Washington, D.C. 20036-5339 Telephone No. (202) 857-6000; Facsimile No. (202) 638-4810

(Status) (patented, pending, abandoned)

(Status) (patented, pending, abandoned)

The undersigned hereby authorizes the U.S. attorneys named herein to accept and follow instructions from the undersigned's assignee, if any, and/or, if the undersigned is not a resident of the United States, the undersigned's domestic attorney, patent attorney or patent agent, as to any action to be take in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and the undersigned. In the event of a change in the person(s) from whom instructions may be taken, the U.S. attorneys named herein will be so notified by the undersigned

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

	Full name of sole or first inventor Francis X. CUNNINGHAM, Jr. Inventor's signature	
	I	Date
	Residence 2727 Washington Avenue, Chevy Chase, MD 20815	
	Citizenship American	
_	Post Office Address Same as the above	
7.00	Full name of sole or second inventor Zairen SUN	
1	Inventor's signature 3 The VI	0/2001
		Date /
	Residence 3405 Tulane Drive, #22, Hyattsville, MD 20783 1083 COPPERSTONE CH	· 0
	Citizenship permant resident, Rockville, MD 20852	$-k \kappa a$
	Post Office Address Same as the above	
	Chinese, P.R. china	
	Full name of sole or third inventor	
	Inventor's signature	
		Date
	Residence	Jule
	Citizenship	
	Post Office Address	
	1 ost Office Address	

ARENT FOX KINTNER PLOTKIN & KAHN, PLLC

Nikaido, Marmelstein, Murray & Oram Intellectual Property Group

Declaration For U.S. Patent Application As a below named inventor, I hereby declare that:

My resi I believ names a	dence, post of e I am the of the listed belo	office address and citic riginal, first and sole w) of the subject man tes of Carotence	zenship are as state e inventor (if only tter which is claime	one name is lised and for which	ited below) or an a patent is soug	tht on the inve	ntion entitl	ed
the spec	ification of w	which is attached here	to unless the follow	ving box is chec	ked:			
\boxtimes	was filed o			s amended on	As PCT Intern	ational Applic	ation	
and/or	was filed o		000	s amended on	As U.S. Paten	t Application		
by any a I acknow I hereby certificately below a	amendment rowledge the du wledge the du y claim forei ate, or §365(and have also	have reviewed and useferred to above. The property to disclose informing priority benefits to any PCT Interpretation identified below any that of the applications.	ation which is mate under 35 U.S.C. § national application foreign application	erial to patentab 119(a)-(d) or §3 which designa for patent or in	ility as defined in 365(b) of any fo ted at least one	n 37 C.F.R. § preign applicat country other	1.56. ion(s) for j than the U ternational	patent or inventor's Juited States, listed Application having
		 					Priority C	laimed No
(List) foreig		mber)	(Country)		/Month/Year Fi		☐ Yes	☐ No
2 2 2 2 2 2 2 2 2	(Nu	mber)	(Country)	(Day	/Month/Year Fi	led)	☐ Yes	□ No
1	(Nu	mber)	(Country)	(Day	/Month/Year Fi	led)	_	_
	claim the be	(Application Num	ber)	Inited States pro (Filing (Filing	Date)	ion(s) listed be	elow.	
		_	ched list for addition		•	applications.		
I hereby designated disclosed duty to	ting the United in the prior disclose information in the prior disclose information in the prior	enefit under 35 U.S. ed States of America application(s) (U.S. rmation which is mat dication and the nation	listed below and, or PCT) in the material to patentability	insofar as the s inner provided t ty as defined in	ubject matter of by the first parag 37 C.F.R. §1.5	each of the claraph of 35, U 6 which became	aims of the S.C. §112	is application is not . I acknowledge the
(List prior Application PCT Inter application designation	ons or national	09/088,724 (Application Serial 09/088,725 (Application Serial	· · · · · · · · · · · · · · · · · · ·	June 2, 1998 (Filing Date) June 2, 1998 (Filing Date)		· · ·		ng, abandoned)
22,980; David T Reg. No Hill, Re and Har	Charles M. Nikaido, I J. 714; M G. 37,33 S. J. Crosby,	the firm of Arent F. Marmelstein, Reg. No. 22,663; Ri furat Ozgu, Reg. No. 51; Carl Schaukowito Reg. No. 44,634.	No. 25,895; George chard J. Berman, 44,275; Robert K ch, Reg. No. 29,2	e E. Oram, Jr., Reg. No. 39,10 Carpenter, Re 11; Kevin Turne	Reg. No. 27,93 7; King L. Wor g. No. 34,794;	1; Douglas H ng, Reg. No. Gregory B. K 437; Rhonda	Goldhush 37,500; Jai ang, Reg. I C. Barton,	, Reg. No. 33,125; mes A. Poulos, III, No. 45,273; Rustan
1 icase C	meet all colli	manications to the 10	nowing address.	1050 Connectic Washington, D	ut Avenue, N.W.C. 20036-5339 (202) 857-6000;	7., Suite 600		3-4810

The undersigned hereby authorizes the U.S. attorneys named herein to accept and follow instructions from the undersigned's assignee, if any, and/or, if the undersigned is not a resident of the United States, the undersigned's domestic attorney, patent attorney or patent agent, as to any action to be take in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and the undersigned. In the event of a change in the person(s) from whom instructions may be taken, the U.S. attorneys named herein will be so notified by the undersigned.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Francis X. CUNNINGHAM, Ir.	Feb. 16, 2001
Inventor's signature Francis & Curringhm, fr.	Date
Residence Citizenship American 2727 Washington Avenue, Chevy Chase, MD 20815	
Post Office Address Same as the above	
Full name of sole or second inventor Zairen SUN	
Inventor's signature	Date
Residence 3405 Tulane Drive, #22, Hyattsville, MD 20783	
Citizenship Post Office Address Same as the above	
Full name of sole or third inventor	
Inventor's signature	Date
Residence	
Citizenship	
Post Office Address	